



Elemental Ratios in Soil: Hidden Indicators of Soil Health, Fertility, and Environmental Change

*Seema Kousar¹, Sachin Kumar¹, Rahul Yadav², Shivam Kumar³ and Chandana S⁴

¹Forest College and Research Institute, Mulugu, Telangana

²Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Uttar Pradesh, India

³Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana

⁴Navsari Agriculture University, Navsari, Gujarat

*Corresponding Author's email: seemakousar954@gmail.com

In soil chemical profile assessment, elemental ratios represent the relative proportions of chemical elements rather than their absolute concentrations, providing a more integrative understanding of soil processes. These ratios are extensively used to interpret soil genesis, nutrient availability, weathering intensity, redox dynamics, and anthropogenic disturbances. Since soils are heterogeneous systems influenced by texture, structure, moisture content, and bulk density, absolute elemental concentrations often vary widely even within short spatial scales. Elemental ratios help minimize these dilution and matrix effects, allowing meaningful comparison between soils across horizons, land-use systems, and climatic regions. From a soil science perspective, elemental ratios serve as diagnostic indicators of pedogenic processes such as eluviation–illuviation, organic matter turnover, leaching of base cations, and accumulation of sesquioxides. They also provide insights into biogeochemical cycling of nutrients and trace elements within soil–plant–microbe systems. Consequently, elemental ratios form a critical component of soil fertility assessment, land degradation studies, and sustainable soil management strategies in both natural and managed ecosystems. (Brady & Weil 2016; Sparks 2003; Alloway 2013)

Conceptual Basis of Elemental Ratios in Soils

The distribution and behavior of elemental ratios in soils are fundamentally controlled by the five soil-forming factors (Fig 1): parent material, climate, organisms, relief, and time. Parent material determines the initial elemental composition, while climate-driven processes such as rainfall and temperature regulate weathering intensity and leaching losses. Biological activity, including root uptake and microbial decomposition, further modifies elemental ratios by redistributing nutrients between soil pools. During pedogenesis, mobile elements such as calcium (Ca), magnesium (Mg), sodium (Na), and potassium (K) are progressively leached from the soil profile under humid conditions, whereas relatively immobile elements such as aluminum (Al), iron (Fe), and titanium (Ti) tend to accumulate. This differential mobility leads to systematic changes in elemental ratios with soil depth and degree of weathering. For example, decreasing Ca/Mg ratios indicate base cation depletion, while declining Si/Al ratios reflect silica loss and advanced soil development. Thus, elemental ratios act as sensitive indicators of soil age, chemical transformation, and pedogenic intensity. (Jenny 1994; Buol et al. 2011; Schaetzl & Thompson 2015).




Fig.1 Factors and process affecting elemental ration in soils

Carbon–Nitrogen–Phosphorus (C:N:P) Ratios in Soil Organic Matter

The C:N:P ratio is a cornerstone of soil chemical and biological assessment, reflecting the stoichiometric balance between carbon, nitrogen, and phosphorus in soil organic matter (SOM). Carbon serves as an energy source for soil microorganisms, while nitrogen and phosphorus are essential nutrients regulating microbial growth and enzymatic activity. In mineral soils, a C:N ratio of approximately 10–12 is generally associated with well-decomposed organic matter and efficient nitrogen mineralization, whereas wider ratios (>20) indicate slow decomposition and temporary nitrogen immobilization. Phosphorus dynamics are often assessed through C:P and N:P ratios, particularly in highly weathered tropical soils where phosphorus availability is limited due to fixation by Fe and Al oxides. High N:P ratios typically indicate phosphorus limitation, which constrains microbial activity and plant productivity. Understanding C:N:P ratios is therefore essential for evaluating soil fertility status, nutrient cycling efficiency, and the sustainability of agricultural and forest ecosystems. (Stevenson 1994; Brady & Weil 2016; Paul 2015)

Base Cation Ratios (Ca:Mg:K:Na) and Soil Fertility

Base cation ratios play a crucial role in determining soil fertility, structure, and plant nutrient availability. Calcium is essential for soil aggregation and structural stability, while magnesium and potassium are vital plant nutrients that can interact antagonistically during uptake. The Ca/Mg ratio is particularly important, as excessively low values may result in clay dispersion, compaction, reduced permeability, and poor root growth. Potassium availability is often evaluated using K/Ca and K/Mg ratios, which help diagnose nutrient imbalances in intensively cultivated soils. High potassium levels relative to calcium or magnesium can suppress the uptake of these nutrients, leading to hidden deficiencies. In forestry and plantation systems, prolonged biomass removal can alter base cation ratios,

contributing to soil acidification and long-term fertility decline. Tisdale et al. 2018); Sparks 2003); Havlin et al. 2016)

Weathering and Pedogenic Indices Based on Elemental Ratios

Elemental ratios involving silicate-forming and sesquioxide-forming elements are widely used as indices of weathering intensity and pedogenic evolution. Ratios such as Si/Al and Si/Fe decrease with increasing weathering due to the preferential loss of silica through leaching and the relative enrichment of aluminum and iron oxides. These trends are especially pronounced in tropical and subtropical environments characterized by intense chemical weathering. Highly weathered soils such as Oxisols and Ultisols typically exhibit low Si/Al ratios, reflecting advanced desilication and dominance of kaolinite and sesquioxide minerals. In contrast, younger soils developed on recent alluvium or volcanic material retain higher Si/Al ratios. These elemental ratios are therefore valuable tools in soil classification, paleopedology, and landscape evolution studies. (Birkeland 1999; Schaetzl & Thompson 2015; Buol et al. 2011).

Redox-Sensitive Elemental Ratios in Soils

Redox-sensitive elemental ratios, particularly Fe/Mn, provide important insights into soil drainage conditions, aeration status, and hydromorphic processes. Under well-aerated conditions, iron and manganese remain in oxidized forms and are relatively immobile. However, under waterlogged or anaerobic conditions, manganese is reduced and mobilized more readily than iron, leading to elevated Fe/Mn ratios. Such ratios are commonly used to identify gleyed horizons, seasonal water stagnation, and wetland soils. In agricultural landscapes, redox-sensitive ratios help assess the suitability of soils for crop production and guide drainage management. They are also critical in understanding the mobility of trace metals and nutrients under fluctuating redox conditions. (Vepraskas & Craft 2016; Sparks 2003; Foth 1991).

Elemental Ratios as Indicators of Soil Pollution

In contaminated soils, elemental ratios are valuable tools for distinguishing natural background levels from anthropogenic inputs. Ratios such as Pb/Zn, Cu/Zn, and Cd/Zn are frequently used to trace pollution sources associated with industrial emissions, mining activities, vehicular traffic, and sewage sludge application. These ratios help overcome limitations associated with absolute concentration data, which may be influenced by soil texture and mineralogy. Normalization against conservative elements such as aluminum or iron further improves interpretation by accounting for natural variability in soil composition. As a result, elemental ratios are widely applied in soil pollution assessment, risk evaluation, and environmental monitoring programs. (Alloway (2013; Kabata-Pendias 2000; Reimann & de Caritat 2005).

Depth-wise Variation of Elemental Ratios in Soil Profiles

Elemental ratios exhibit systematic variation with soil depth due to pedogenic processes such as eluviation and illuviation. Surface horizons are typically enriched in organic carbon, nitrogen, and phosphorus, resulting in narrower C:N ratios compared to deeper layers. With increasing depth, organic matter content declines, and mineral-associated elements such as iron and aluminum become more dominant. Subsoil horizons often show accumulation of Fe and Al oxides, leading to lower Si/Al ratios and reflecting advanced weathering. Analyzing depth-wise trends in elemental ratios provides valuable information on horizon differentiation, nutrient translocation, and long-term soil development. (Schaetzl & Thompson 2015; Brady & Weil 2016; Jenny 1994)

Importance of Elemental Ratios in Sustainable Soil Management

Elemental ratios offer integrative indicators for assessing soil health and guiding sustainable land management practices. They are increasingly used in fertilizer recommendation strategies, soil conservation planning, and land-use suitability assessment. In forestry and

agroforestry systems, elemental ratios help evaluate long-term nutrient depletion caused by biomass harvesting and inform sustainable management interventions. Moreover, elemental ratios play a significant role in climate-smart soil management by linking carbon sequestration potential with nutrient cycling efficiency. Understanding these relationships is essential for enhancing soil resilience, productivity, and ecosystem services under changing climatic conditions. (Lal 2015; Brady & Weil 2016; Paul 2015). From a soil science perspective, elemental ratios are indispensable tools for chemical profile assessment, offering deeper insights into soil fertility, weathering processes, redox dynamics, pollution status, and pedogenic history. When interpreted alongside absolute elemental concentrations, soil morphological observations, and land-use information, elemental ratios significantly enhance the understanding of soil functioning. Their application is therefore central to informed decision-making in agriculture, forestry, environmental management, and sustainable land-use planning. (Sparks (2003); Alloway (2013); Schaetzl & Thompson (2015).

Stoichiometric Relationships and Ecological Balance in Soil

Beyond basic C:N:P evaluation, ecological stoichiometry provides a broader theoretical framework for understanding elemental balance in soil ecosystems. Soil organisms require nutrients in relatively fixed proportions for growth and metabolism. When elemental ratios deviate from optimal microbial requirements, nutrient immobilization or mineralization processes are altered. For instance, if organic residues possess high C:N ratios, microorganisms immobilize available nitrogen to satisfy their metabolic demand, temporarily reducing nitrogen availability for plants. Stoichiometric imbalance also affects greenhouse gas emissions. Excess nitrogen relative to carbon may enhance nitrification and denitrification processes, increasing nitrous oxide emissions. Conversely, balanced C:N ratios promote stable humus formation and long-term carbon sequestration. Therefore, elemental ratios are closely linked to ecosystem productivity, soil respiration, and climate regulation. These relationships are increasingly studied under global climate change scenarios to understand soil feedback mechanisms (Sterner & Elser, 2002; Cleveland & Liptzin, 2007).

Elemental Ratios in Soil Classification and Taxonomy

Elemental ratios play a diagnostic role in modern soil classification systems. In highly weathered soils, low Si/Al and Si/Fe ratios indicate advanced desilication and dominance of sesquioxides. Such chemical signatures help differentiate mature tropical soils from younger temperate soils. Similarly, base saturation ratios derived from Ca, Mg, K, and Na content are important criteria in soil taxonomy for distinguishing between acidic and base-rich soil orders. In saline and sodic soils, Na/Ca and Na/Mg ratios are particularly significant. Elevated sodium relative to calcium leads to dispersion of clay particles, poor aggregation, and reduced permeability. Thus, elemental ratios are not merely chemical indicators but also structural and functional determinants of soil behavior. Integrating chemical ratios with morphological characteristics enhances the precision of soil classification and land capability assessment.

Analytical Methods for Determining Elemental Ratios

Accurate determination of elemental ratios depends on reliable analytical techniques. Soil samples are typically air-dried, ground, and sieved before laboratory analysis. Total elemental concentrations may be determined using methods such as atomic absorption spectrophotometry (AAS), inductively coupled plasma optical emission spectroscopy (ICP-OES), or X-ray fluorescence (XRF). Organic carbon is commonly measured by Walkley–Black wet oxidation or dry combustion methods, while nitrogen is determined using the Kjeldahl method. Once elemental concentrations are obtained, ratios are calculated by dividing the molar or mass-based values of the respective elements. For pedogenic interpretation, molar ratios are often preferred because they reflect atomic relationships more accurately than weight-based ratios. Quality control procedures, including the use of standard reference materials and duplicate analysis, are essential to ensure data reliability. Proper

sampling design and statistical analysis further strengthen the interpretation of elemental ratio data.

Limitations and Precautions in Interpreting Elemental Ratios

Although elemental ratios provide valuable integrative indicators, their interpretation must consider certain limitations. Soil heterogeneity, sampling depth variability, seasonal fluctuations, and land-use history can influence ratio values. For example, recent fertilizer application may temporarily distort natural nutrient ratios. Similarly, erosion or deposition processes may modify surface soil chemistry independent of pedogenic development. Another limitation is that elemental ratios do not always reveal absolute nutrient sufficiency or deficiency. A balanced ratio may still correspond to low overall nutrient availability. Therefore, elemental ratios should be interpreted alongside absolute concentrations, soil pH, cation exchange capacity, texture, and biological indicators. Integrated soil assessment approaches ensure more accurate conclusions.

Role of Elemental Ratios in Precision Agriculture and Modern Soil Management

With the advancement of precision agriculture, elemental ratios are increasingly used to guide site-specific nutrient management. Spatial mapping of Ca/Mg or N/P ratios using geospatial tools enables targeted fertilizer application, improving nutrient use efficiency and reducing environmental losses. Variable rate technology allows farmers to correct nutrient imbalances based on ratio analysis, thereby enhancing crop productivity while minimizing costs. In sustainable forestry and agroecosystems, monitoring long-term trends in elemental ratios helps evaluate nutrient depletion due to biomass harvesting. Restoration programs often use elemental ratio data to design balanced fertilization strategies that promote soil recovery. Thus, elemental ratios contribute significantly to sustainable intensification and resource-efficient land management.

Conclusion

Elemental ratios represent powerful diagnostic tools in soil chemical profile assessment. They integrate complex pedogenic, biological, and environmental processes into interpretable indicators of soil functioning. From nutrient cycling and fertility evaluation to pollution monitoring and soil classification, elemental ratios provide a comprehensive framework for understanding soil systems. When combined with modern analytical techniques and geospatial technologies, elemental ratio analysis supports informed decision-making in agriculture, forestry, environmental monitoring, and climate adaptation strategies. Therefore, the study of elemental ratios remains central to contemporary soil science and sustainable land management.

References

1. Alloway, B. J. (2013). Trace metals and metalloids in soils and their bioavailability. *Heavy Metals in Soils*, 22, 4-18.
2. Birkeland, P. W. (1999). *Soils and geomorphology* (3rd ed.). Oxford University Press.
3. Brady, N. C., & Weil, R. R. (2016). *The Nature and Properties of Soils* 15th edition New York.
4. Buol, S. W., Southard, R. J., Graham, R. C., & McDaniel, P. A. (2011). *Soil genesis and classification*. John Wiley & Sons.
5. Cleveland, C. C., & Liptzin, D. (2007). C: N: P stoichiometry in soil: is there a "Redfield ratio" for the microbial biomass?. *Biogeochemistry*, 85(3), 235-252.
6. Foth, H. D. (1991). *Fundamentals of soil science* (No. Ed. 8, pp. xv+-360).
7. Havlin, J. L., Tisdale, S. L., Nelson, W. L., & Beaton, J. D. (2016). *Soil fertility and fertilizers*. Pearson Education India.
8. Jenny, H. (1994). *Factors of soil formation: a system of quantitative pedology*. Courier Corporation.
9. Kabata-Pendias, A. (2000). *Trace elements in soils and plants*. CRC press.

10. Lal, R. (2015). Restoring soil quality to mitigate soil degradation. *Sustainability*, 7(5), 5875-5895.
11. Paul, E. A. (2015). Soil microbiology, ecology, and biochemistry: an exciting present and great future built on basic knowledge and unifying concepts. *Soil Microbiology, Ecology and Biochemistry, fourth ed. Academic Press San Diego*, 1-15.
12. Reimann, C., & de Caritat, P. (2005). Distinguishing between natural and anthropogenic sources for elements in the environment: regional geochemical surveys versus enrichment factors. *Science of the total environment*, 337(1-3), 91-107.
13. Schaetzl, R. J., & Thompson, M. L. (2015). *Soils*. Cambridge university press.
14. Sparks, D. L. Environmental Soil Chemistry: 2003. *Chemistry*, 6(7), 8.
15. Sterner, R. W., & Elser, J. J. (2003). *Ecological stoichiometry: the biology of elements from molecules to the biosphere*. Princeton university press.
16. Stevenson, F. J. (1994). *Humus chemistry: genesis, composition, reactions*. John Wiley & Sons.
17. Tisdale, S. L., Nelson, W. L., Havlin, J. L., & Beaton, J. D. (2018). *Soil fertility and fertilizers* (8th ed.). Pearson.
18. Vepraskas, M. J., & Craft, C. B. (Eds.). (2016). *Wetland soils: genesis, hydrology, landscapes, and classification*. CRC press.