

Increasing Boll Number and Boll Size through Efficient Management of Sympodial Growth in Cotton

*Vijay Sundar B, Dr. Suchismita Tripathy, Dr. Jyoti Rekha Patnaik and Krishnakumar S

College of Agriculture, OUAT, Bhubaneswar, Odisha, India

*Corresponding Author's email: ysundar493@gmail.com

Cotton (*Gossypium hirsutum* L.) holds a preeminent position in global agriculture as the primary source of natural textile fiber and a significant contributor to edible oil and seed cake. The unique botanical characteristic of cotton is its indeterminate growth habit, which allows it to simultaneously undergo vegetative growth and reproductive development over a prolonged period. The ultimate economic yield of a cotton crop is mathematically defined by the product of plant population per hectare, the number of harvested bolls per plant, and the average weight of those bolls. Among these yield components, boll number and boll weight are the most critical and highly malleable traits, fundamentally governed by the architecture and physiology of the plant. Specifically, these reproductive units are borne exclusively on sympodial (fruiting) branches. The efficient management of sympodial growth—ensuring an optimal rate of branch initiation, maximizing floral bud (square) retention, and providing adequate assimilates for boll filling—is the cornerstone of modern cotton agronomy. This document explores the advanced crop physiology of sympodial growth and outlines practical agronomic management strategies to optimize the critical balance between boll number and boll size.

Cotton Plant Architecture

To manage cotton efficiently, a thorough understanding of its dimorphic branching pattern is essential. The cotton plant exhibits two distinct types of branches: monopodial and sympodial. Monopodial branches (vegetative branches) typically emerge from axillary buds located at the lower nodes of the main stem (nodes 1 through 5, depending on the variety and environment). They grow at an acute angle to the main stem, exhibiting continuous, straight apical growth. Monopodia do not bear fruit directly; instead, they act similarly to the main stem, eventually producing secondary sympodial branches. Sympodial branches (fruiting branches) emerge from extra-axillary buds at higher nodes (usually starting at node 6 or 7). These branches exhibit a distinctive zigzag morphological pattern. The terminal meristem of a sympodium produces a floral bud (square), and the branch continues to grow via a lateral bud, resulting in a series of nodes, each bearing a leaf and a fruit.




Figure 1. Schematic representation of monopodial and sympodial branching in cotton.

Characteristic	Monopodial Branches	Sympodial Branches
Growth Habit	Vegetative, straight, upward	Reproductive, zigzag, horizontal
Origin	Axillary buds, lower nodes	Extra-axillary buds, upper nodes
Fruiting Mechanism	Indirectly bears bolls on secondary branches	Directly bears squares/bolls at each node
Terminal Bud	Continues vegetative growth	Terminates in a floral structure (square)

Physiology of Sympodial Growth

The initiation and development of sympodial branches dictate the yield potential of the plant. A new sympodial node (containing a square and a subtending leaf) is produced approximately every 3 to 6 days, highly influenced by temperature and thermal time (degree days).




Figure 2. Typical developmental timeline of a fruiting form on a sympodial branch.

Source-Sink Relationships and Assimilate Partitioning: In the context of cotton physiology, mature green leaves act as carbohydrate 'sources', while active meristems, roots, and developing bolls act as 'sinks'. On a sympodial branch, the subtending leaf associated with a specific boll is the primary source of assimilates for that fruit. During boll development, the subtending leaf provides nearly 60-87% of the required photosynthate (Mangi et al., 2021). If the subtending leaf is damaged, shaded, or senesces prematurely, the developing boll must rely on assimilates translocated from main stem leaves or other sympodia, which severely limits its final size.




Figure 3. Assimilate flow from subtending leaf (source) to developing boll (sink).

Boll Number Determination

While a cotton plant can initiate a vast number of squares, only a fraction survive to become mature harvestable bolls. The final boll number is mathematically determined by the rate of

node addition minus the rate of abscission (shedding). Natural shedding of 40–60% of fruiting forms is common. Shedding is a highly regulated physiological mechanism designed to adjust the plant's reproductive sink load to match its carbohydrate source capacity. **Factors Controlling Shedding:** Shedding is primarily hormonally driven. Stresses such as water deficit, nutrient starvation, low light (cloudy days or dense canopy shading), or extreme heat trigger an increase in ethylene and abscisic acid (ABA) synthesis in the young bolls, accompanied by a drop in indole-3-acetic acid (IAA/auxin). This hormonal imbalance activates hydrolytic enzymes (cellulase and polygalacturonase) which degrade cell walls in the abscission zone at the base of the pedicel, resulting in fruit drop.

Physiological Pathway of Boll Shedding




Figure 4. Physiological mechanism of square and boll shedding in response to stress.

Boll Size/Weight Determination

Boll size and weight are determined by two main physiological components: the number of fertilized seeds per boll and the mass of lint fibers produced per seed. The development of a cotton boll occurs in three overlapping phases:

1. Enlargement Phase (0–20 days post-anthesis): Rapid cell division and elongation occur. Water relations are paramount here; water deficit reduces cell turgor, restricting final boll volume.
2. Filling Phase (20–45 days post-anthesis): Characterized by secondary wall thickening and massive cellulose deposition in the fiber cells. This phase is highly energy-intensive and depends entirely on the continuous supply of sucrose from the subtending leaf.
3. Maturation Phase: Drying and boll opening.

Nutrient Requirements for Boll Size: Potassium (K) is arguably the most critical macronutrient during the boll-filling phase. K acts as the primary osmoticum required to maintain turgor pressure for fiber elongation. Furthermore, K is strictly required for the efficient phloem loading and transport of sucrose from the source leaves into the developing bolls. Potassium deficiency during this phase invariably leads to smaller bolls, premature leaf senescence, and reduced fiber micronaire.

Efficient Management of Sympodial Growth

Agronomic interventions aim to optimize the initiation, retention, and filling of bolls on sympodial branches. Key management practices include:

- **Variety and Plant Architecture:** Modern, compact Bt cotton hybrids with early maturity tend to partition a higher percentage of biomass into reproductive structures (higher Harvest Index) compared to traditional leafy varieties.
- **Plant Density and Spacing:** High-Density Planting Systems (HDPS) are increasingly adopted. While HDPS reduces the number of sympodia and bolls per individual plant, it significantly increases the total number of bolls per hectare, compensating for individual plant yield reductions.
- **Sowing Time:** Timely sowing ensures that the critical squaring and flowering phases coincide with optimal temperatures and maximum solar radiation, enhancing photosynthetic rates of subtending leaves.
- **Nutrient Management:** Nitrogen (N) drives vegetative growth and leaf area index (LAI). Over-application of N causes excessive vegetative (rank) growth, shading of lower sympodia, and increased shedding. Phosphorus (P) is vital for energy transfer during flowering. Potassium (K) application (often through foliar sprays of KNO₃ at boll development) is essential to maximize boll weight.
- **Irrigation and Moisture:** Maintaining soil moisture near field capacity during peak flowering and early boll development is critical to prevent ethylene-induced shedding and to maintain cell turgor for fiber elongation.

Plant Growth Regulators (Mepiquat Chloride)

Canopy management often relies on the application of Plant Growth Regulators (PGRs) like mepiquat chloride (MC). MC is a systemic anti-gibberellin that inhibits cell elongation. Physiologically, it reduces internode length on both the main stem and sympodia, resulting in a more compact canopy that improves light penetration. This limits excessive vegetative sink activity, redirecting assimilates toward the developing bolls, thereby improving retention and weight.




Figure 5. Relative physiological response of cotton to Mepiquat Chloride application.

Topping

Topping involves the removal of the apical meristem (main stem apex) to arrest terminal vegetative growth. It eliminates apical dominance, forcing the plant to channel photoassimilates entirely into the existing sympodial bolls. While manual topping is labor-intensive, recent studies emphasize chemical topping. Qi et al. (2023) demonstrated that chemical topping with mepiquat chloride during the flowering period suppresses apical dominance without compromising maturity or spatial boll distribution.

Optimum Vegetative-Reproductive Balance

The ultimate goal of sympodial management is to maintain a harmonious vegetative-reproductive balance. Excessive vegetative growth (rank growth) acts as a massive sink, starving developing bolls of carbohydrates. Conversely, if the plant shifts too early into heavy reproductive growth (premature cutout), it lacks the leaf area (source) required to fill the bolls, leading to small, low-quality fruit. Achieving the optimum balance requires synchronized use of N-fertilization, strategic irrigation deficits prior to flowering, and timely MC applications.

Critical Growth Stages for Management

Different agronomic inputs must be timed to match the physiological demands of the crop.

Growth Stage	Physiological Event	Key Management Focus
Squaring (35-45 DAS)	Floral initiation, initial sympodial growth	Weed control, initial N application, prevent moisture stress
Flowering (60-80 DAS)	Pollination, maximum sink formation	Mepiquat chloride application, peak irrigation, pest monitoring
Boll Setting (80-100 DAS)	Cell elongation in fibers, high risk of shedding	Maintain turgor pressure (water), ensure N & P availability
Boll Filling (100-130 DAS)	Cellulose deposition, maximum assimilate demand	Foliar Potassium (KNO_3), chemical topping to stop vegetative growth

Trade-off Between Boll Number and Boll Size

Agronomists frequently encounter a negative correlation between the number of bolls per plant and the average weight per boll. As the number of retained bolls increases, the competition for a finite pool of carbohydrates intensifies. Simply pushing a crop to retain maximum bolls via excessive nitrogen or PGRs can result in a high population of undersized bolls with poor fiber micronaire. Successful agronomy identifies the 'sweet spot' on this trade-off curve where yield is maximized without compromising fiber quality.




Figure 6. The physiological trade-off between boll number and boll weight due to assimilate competition.

Integrated Strategy

Achieving high yields requires an integrated approach that connects management practices directly to physiological outcomes.




Figure 7. Flowchart mapping management inputs to physiological responses and yield.

Important Findings from Recent Research (2020–2026)

Recent literature has provided deeper insights into the source-sink dynamics and advanced canopy management techniques in cotton.

- **Source-Sink Dissection:** Mangi et al. (2021) rigorously quantified the importance of the subtending leaf in upland cotton. By dissecting the source-sink relationship, they confirmed that the subtending leaf is the primary driver of fiber quality and yield attributes, providing the vast majority of assimilates directly to the associated boll.
- **Chemical Topping Innovations:** Traditional manual topping is highly effective but economically unviable due to labor constraints. Research by Qi et al. (2023) demonstrated that chemical topping (CT) utilizing high doses of mepiquat chloride at flowering acts as a perfect substitute. CT successfully regulated the spatial distribution of bolls and controlled apical dominance without negatively impacting the accumulation of boll fractions, ensuring that the crop matures efficiently and maintains high yield potential.

Practical Recommendations & Key Take-Home Points

To translate these physiological principles into field-level success, agronomists and producers should adhere to the following recommendations:

1. Adopt High-Density Planting Systems (HDPS) combined with compact varieties to shift the reliance from individual plant performance to population performance.
2. Monitor the node above white flower (NAWF) to accurately track the transition from vegetative to reproductive growth and anticipate physiological cutout.
3. Apply nitrogen in splits to avoid rank growth during the pre-flowering stage, saving the bulk for peak reproductive demand.
4. Prioritize Potassium (K) nutrition. Utilize foliar sprays of 2% KNO₃ during the boll filling stage to sustain fiber elongation and sucrose transport.
5. Utilize Mepiquat Chloride (MC) proactively rather than reactively. Apply low doses at early squaring and increase rates as the canopy closes.
6. Implement chemical topping in regions with labor shortages to arrest terminal growth and redirect energy to late-season bolls.

7. Protect the subtending leaves at all costs. Late-season insect management should prioritize preserving the canopy that directly feeds the developing bolls.
8. Prevent moisture stress particularly between 60 to 100 days after sowing to minimize ethylene-induced square and boll shedding.
9. Acknowledge the trade-off curve: Do not attempt to force retention of late-season squares if the plant lacks the thermal time or leaf area to mature them into heavy bolls.
10. Constantly balance the 'Source' (leaf area index and photosynthetic efficiency) with the 'Sink' (boll load) to achieve sustainable, high-quality lint yields.

Conclusion

Increasing boll number and boll size is not achieved through a single input, but through the holistic management of the cotton plant's sympodial growth. By understanding the physiological triggers of branch initiation, abscission, and assimilate partitioning, agronomists can utilize spacing, precise fertilization, and PGRs to architect a canopy optimized for high yield and superior fiber quality.

References

1. Mangi, N., Nazir, M. F., Wang, X., Iqbal, M. S., Sarfraz, Z., Jatoi, G. H., Mahmood, T., Ma, Q., & Shuli, F. (2021). Dissecting source-sink relationship of subtending leaf for yield and fiber quality attributes in upland cotton (*Gossypium hirsutum* L.). *Plants*, 10(6), 1147. <https://doi.org/10.3390/plants10061147>
2. Qi, H., Xiao, C., Zhao, W., Xu, D., Eneji, A. E., Lu, Z., Shao, R., Wang, G., Du, M., Tian, X., & Li, Z. (2023). Chemical topping with mepiquat chloride at flowering does not compromise the maturity or yield of cotton. *Agronomy*, 13(2), 497. <https://doi.org/10.3390/agronomy13020497>