

Genetic Improvement of Red Cabbage: Opportunities for Nutrition, Productivity and Resilience

Veer Bhan Singh¹, Dr. Muzeev Ahmad¹, *Manasi Singh¹ and Yatharth Mishra²

¹Department of Agriculture, IAST, Integral University, Lucknow, Uttar Pradesh, India

²Department of Genetics and Plant Breeding, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, Uttar Pradesh, India

*Corresponding Author's email: manasisingh112001@gmail.com

Red cabbage is more than a colourful vegetable. Its unusual phytochemical profile, expanding genomic resources and exploitable genetic diversity make it a strong candidate for next-generation breeding aimed at healthier diets, dependable yield and climate resilience. The breeding challenge is no longer “yield or quality.” The opportunity is to design red cabbage for nutrition, marketable yield and resilience at the same time.

Introduction

Red cabbage (*Brassica oleracea* var. *capitata* f. *rubra*) combines the familiar agronomic value of heading cabbage with a distinctive biochemical trait: the accumulation of red-to-purple anthocyanin pigments. Recent reviews confirm that red cabbage cultivars are especially rich in phenolics and anthocyanins, while their glucosinolate composition varies substantially among cultivars and environments. This makes red cabbage unusual from a breeder's perspective: colour, nutritional quality, processing value and market appeal are biologically linked, yet each can still be improved independently. The central opportunity is therefore not simply to breed a darker head. Modern improvement can target a package of traits—stable pigment composition, desirable glucosinolates, vitamin C, head yield and compactness, uniform maturity, resistance to major diseases, and tolerance to heat and water stress. The availability of high-quality *Brassica oleracea* reference genomes and a recent graph-based pan-genome now gives breeders a much richer map of the structural and allelic variation behind these traits.

Nutrition: breeding the chemistry of colour

Anthocyanins are the signature compounds of red cabbage. They are dominated by cyanidin derivatives, many of which are acylated. Acylation improves pigment stability, a valuable feature for fresh-market appearance and for the use of red-cabbage extracts as natural food colourants. When 27 red cabbage genotypes were profiled across two growing seasons, researchers detected 29 anthocyanins and three distinct profile “fingerprints”, demonstrating that breeders have substantial biochemical diversity to work with [2]. The practical implication is important: selection can focus not only on total pigment concentration but also on the composition and stability of the pigment profile. The genetics of purple pigmentation are becoming increasingly tractable. Loss of expression of BoMYBL2-1, a negative regulator of anthocyanin biosynthesis, was shown to underline purple coloration in cabbage, and molecular markers were validated across diverse cabbage lines. A subsequent genome-wide study identified 88 anthocyanin-biosynthetic genes in *B. oleracea* and highlighted BoMYB113.1 as a positive regulator and BoMYBL2.1 as a negative regulator of anthocyanin accumulation. These findings create a route from visual selection to marker-assisted selection, particularly where breeders want consistent pigmentation across seasons or wish to combine high anthocyanin content with other commercial traits.



Figure 1- represents the anthocyanin pigment in red cabbage

Red cabbage also offers opportunities beyond anthocyanins. Glucosinolates are central Brassica defence metabolites and important targets in nutritional breeding. Screening of 146 cabbage genotypes revealed large variation in individual and total glucosinolate classes, including glucoerucin, sinigrin and glucoraphanin. This diversity means nutritional improvement can be based on real genetic variation rather than on agronomic manipulation alone. However, “more” is not automatically “better”: glucosinolate profiles influence flavour, pungency and plant defence, so breeding goals should balance nutritional composition with consumer acceptance and crop performance.




Figure 1. An integrated breeding roadmap for red cabbage. Nutrition, productivity and resilience are treated as simultaneous selection targets, supported by germplasm diversity, precise phenotyping, hybrid breeding, markers, genomics and gene editing. Synthesis based on

Productivity: build nutrition into a better head

Nutritional quality has limited value if the cultivar is low-yielding, non-uniform or poorly adapted. Red cabbage breeding must therefore treat head weight, compactness, core length, maturity, field uniformity and resistance to splitting as primary economic traits. A 2024 evaluation of 28 red cabbage genotypes in the north-western Himalayas found meaningful genetic variability for net head weight, stalk and core length, non-wrapper leaves and vitamin C content; the most divergent clusters were proposed as useful parents for creating superior recombinants and hybrids [6]. Such studies are especially valuable because they identify

diversity directly within red cabbage rather than extrapolating entirely from white cabbage or other *B. oleracea* crops. Hybrid breeding offers a powerful productivity route. In cabbage, cytoplasmic male sterility (CMS) simplifies F1 seed production, while doubled haploid (DH) technology rapidly creates homozygous parental lines. Indian research using Ogura-CMS and DH cabbage parents identified several F1 combinations that out-yielded a commercial check while maintaining strong uniformity [9]. Newer haploid-induction systems may accelerate this pipeline further: a BoCENH3-based system was recently used to generate homozygous fertile and CMS lines in a single breeding framework. For red cabbage, these tools could shorten the time needed to combine pigment quality with heterosis for head yield and market uniformity. Genomics is also beginning to clarify the architecture of heading itself. The 2020 D134 cabbage reference genome assembled a 529.9-Mb chromosome-level genome with 44,701 annotated protein-coding genes. The newer *B. oleracea* pan-genome captures structural variants that a single reference can miss and has identified genes associated with leaf and heading diversification. This matters because selection for “good head type” is selection on a complex developmental phenotype—leaf curvature, packing, timing, architecture and source–sink balance—and structural variation may explain useful differences that are invisible to conventional marker sets.

Resilience: protect colour and yield under stress

Red cabbage is a cool-season crop, so rising temperatures threaten both head formation and quality. Experiments comparing white and red cabbage cultivars under heat stress showed clear differences in membrane thermostability, chlorophyll retention and head formation. Expression of BoHSP70, BoSCL13 and BoDPB3-1 also differentiated heat-tolerant from heat-sensitive material, providing potential physiological and molecular selection indicators. For breeders, the message is straightforward: heat tolerance should be measured at the head-forming stage and linked to marketable yield, not only to seedling survival. Disease resilience is equally important. Black rot, caused by *Xanthomonas campestris* pv. *campestris*, can be devastating in cabbage production. Screening of 162 cabbage accessions found only four highly resistant sources to the tested race, while inheritance analysis indicated contributions from major genes and polygenes. Clubroot, caused by *Plasmodiophora brassicae*, is another difficult soil-borne disease. QTL-sequencing in *B. oleracea* identified three resistance QTL on chromosome C07 and a functional marker with about 95% accuracy in the tested population. Such resistance loci are ideal candidates for introgression into red cabbage backgrounds because host resistance reduces reliance on chemical or soil-based disease management. Marker-assisted breeding can now address several cabbage traits simultaneously. A recent review catalogued markers associated with CMS, male sterility, Fusarium wilt, black rot, clubroot, downy mildew, head splitting, bolting and temperature responses, while also emphasizing that complex polygenic traits remain less well served by validated markers. This is where genomic selection, high-throughput phenotyping and multi-environment trials can add value—especially for resilience traits that are strongly influenced by genotype $\times$ environment interaction.

From single traits to multi-trait design

The next phase of red cabbage improvement should move beyond sequential breeding—first yield, then colour, then disease resistance—and instead use multi-trait selection from the beginning. A practical breeding population can be screened simultaneously for anthocyanin profile, glucosinolate composition, vitamin C, head yield, compactness, splitting, maturity and stress response. Parents can then be selected using genomic relationships, trait-linked markers and combining ability, while DH and CMS technologies speed the production of uniform hybrids. Genome editing may eventually provide precise options for validated genes, but it should complement rather than replace germplasm-based breeding. The strongest long-term strategy is to preserve and characterize broad red-cabbage diversity, use pangenomic information to identify useful alleles and structural variants, and deploy precision tools only where gene function is clear. This approach also reduces the risk of narrowing the crop's genetic base while chasing a few high-profile traits.

For growers and consumers, the ideal outcome is simple even if the genetics are complex: a red cabbage that forms a uniform, dense head; retains an attractive, stable colour; delivers a favourable spectrum of health-related phytochemicals; tolerates warmer and more variable seasons; and resists major diseases without sacrificing yield. With red-cabbage-specific diversity now being documented and the *B. oleracea* genomic toolbox expanding rapidly, that goal is increasingly realistic rather than aspirational.

Breeding priorities immediately

Nutrition	Stable, high-value anthocyanin profiles; favourable glucosinolate balance; vitamin C; colour retention and sensory quality.
Productivity	High net head weight, compactness, uniform maturity, reduced core, resistance to splitting and strong F1 performance.
Resilience	Heat tolerance during heading, stable yield across environments, and durable resistance to black rot and clubroot.

Conclusion

Red cabbage possesses considerable genetic and biochemical diversity that can be exploited for cultivar improvement. Variation in anthocyanins, glucosinolates and vitamin C provides measurable targets for nutritional breeding, while molecular studies are beginning to reveal genes underlying anthocyanin accumulation. The next generation of red-cabbage breeding should integrate germplasm diversity, precise phenotyping, biochemical analysis, molecular markers, doubled haploids, hybrid breeding and multi-environment testing. The goal is simple: “Breed red cabbage not only for colour, but for yield, nutrition, stability and resilience.”

References

1. Statilko, O., Tsiaka, T., Sinanoglou, V. J., & Strati, I. F. (2024). Overview of phytochemical composition of *Brassica oleracea* var. *capitata* cultivars. *Foods*, 13, 3395. <https://doi.org/10.3390/foods13213395>
2. Strauch, R. C., Mengist, M. F., Pan, K., Yousef, G. G., Iorizzo, M., Brown, A. F., & Lila, M. A. (2019). Variation in anthocyanin profiles of 27 genotypes of red cabbage over two growing seasons. *Food Chemistry*, 301, 125289. <https://doi.org/10.1016/j.foodchem.2019.125289>
3. Song, H., Yi, H., Lee, M., Han, C.-T., Lee, J., Kim, H., Park, J.-I., Nou, I., Kim, S.-J., & Hur, Y. (2018). Purple *Brassica oleracea* var. *capitata* f. *rubra* is due to the loss of BoMYBL2-1 expression. *BMC Plant Biology*, 18. <https://doi.org/10.1186/s12870-018-1290-9>
4. Han, F., Zhang, X., Yang, L., Zhuang, M., Zhang, Y., Liu, Y., Li, Z., Wang, Y., Fang, Z., Ji, J., & Lv, H. (2021). Genome-wide characterization and analysis of the anthocyanin biosynthetic genes in *Brassica oleracea*. *Planta*, 254. <https://doi.org/10.1007/s00425-021-03746-6>
5. Bhandari, S. R., Rhee, J., Choi, C. S., Jo, J. S., Shin, Y. K., & Lee, J. G. (2020). Profiling of individual desulfo-glucosinolate content in cabbage head germplasm. *Molecules*, 25, 1860. <https://doi.org/10.3390/molecules25081860>
6. Deep, N., Bhardwaj, R. K., Kumari, R., & Bhan, S. (2024). Variability and genetic divergence studies in red cabbage under North-Western Himalayas. *Vegetable Science*, 51(1). <https://doi.org/10.61180/vegsci.2024.v51.i1.10>
7. Lv, H., Wang, Y., Han, F., Ji, J., Fang, Z., Zhuang, M., Li, Z., Zhang, Y., & Yang, L. (2020). A high-quality reference genome for cabbage obtained with SMRT reveals novel genomic features and evolutionary characteristics. *Scientific Reports*, 10. <https://doi.org/10.1038/s41598-020-69389-x>
8. Guo, N., Wang, S., Wang, T., et al. (2024). A graph-based pan-genome of *Brassica oleracea* provides new insights into its domestication and morphotype diversification. *Plant Communications*, 5. <https://doi.org/10.1016/j.xplc.2023.100791>

9. Aruna, T. S., Parkash, C., Kumar, S., Tomar, B. S., Behera, T. K., Singh, R., Singh, K., & Kumar, S. (2023). Utilizing Ogura-CMS and doubled haploid lines of cabbage for new-generation hybrid breeding. *Vegetable Science*, 50(1). <https://doi.org/10.61180/vegsci.2023.v50.i1.02>
10. Strembovskiy, I. V., & Kroupin, P. (2025). The current status and prospects of molecular marker applications in head cabbage: A review. *Agronomy*, 15, 2644. <https://doi.org/10.3390/agronomy15112644>
11. Kong, C., Chen, G., Yang, L., Zhuang, M., Zhang, Y., Wang, Y., Ji, J., Fang, Z., & Lv, H. (2021). Germplasm screening and inheritance analysis of resistance to cabbage black rot in a worldwide collection of cabbage resources. *Scientia Horticulturae*, 110234. <https://doi.org/10.1016/j.scienta.2021.110234>
12. Ce, F., Mei, J., He, H., Zhao, Y., Hu, W., Yu, F., Li, Q., Ren, X., Si, J., Song, H., & Qian, W. (2021). Identification of candidate genes for clubroot-resistance in *Brassica oleracea* using QTL-sequencing. *Frontiers in Plant Science*, 12, 703520. <https://doi.org/10.3389/fpls.2021.703520>
13. Moradpour, M., Abdullah, S., & Namasivayam, P. (2021). The impact of heat stress on morpho-physiological response and expression of specific genes in *Brassica oleracea*. *Plants*, 10, 1064. <https://doi.org/10.3390/plants10061064>
14. Han, F., Zhang, X., Liu, Y., Liu, Y., Zhao, H., & Li, Z. (2024). One-step creation of CMS lines using a BoCENH3-based haploid induction system in *Brassica* crop. *Nature Plants*, 10, 581–586. <https://doi.org/10.1038/s41477-024-01643-w>