

Optimizing Sowing Time and Sowing Management in Hybrid Sunflower for Higher Seed and Oil Yield

*Krishnakumar S, Dr. Suchismita Tripathy, Dr. Jyoti Rekha Patnaik and Vijay Sundar B

College of Agriculture, OUAT, Bhubaneswar, Odisha, India

*Corresponding Author's email: krishnakumarsenthil2004@gmail.com

Sunflower (*Helianthus annuus* L.) is one of the most important oilseed crops globally, prized for its high-quality edible oil rich in polyunsaturated fatty acids (PUFA), particularly linoleic and oleic acids. With the advent of high-yielding hybrids, sunflower cultivation has shifted from marginal subsistence to highly commercial farming. Unlike traditional varieties, modern hybrids are highly responsive to management practices and environmental variables. Among these, sowing time is the most critical non-monetary input. Sowing date dictates the environmental conditions (temperature, photoperiod, solar radiation, and moisture) that the crop will experience at various phenological stages, ultimately determining biological productivity, harvest index, seed yield, and oil accumulation.

Why Sowing Optimization Matters

Optimizing the sowing date aligns the crop's physiological demands with the optimal climatic windows. Genotype $\times$ Environment $\times$ Management ($G \times E \times M$) interactions heavily influence sunflower performance. Sowing too early may expose seedlings to low temperatures or frost, slowing germination and reducing uniform establishment. Conversely, delayed sowing forces the critical reproductive phases (flowering and seed filling) into periods of heat stress or moisture deficit, shortening crop duration and drastically reducing seed weight and oil yield.




Figure 1: Sowing Date to Yield Pathway

Table 1: Response Matrix of Sowing Windows

Sowing Window	Temperature / Weather	Growth Response	Seed Yield	Oil Content	Oil Yield
Early Sowing	Cooler initial, increasing later	Slow emergence, extended vegetative phase	Moderate to High	High (if no terminal heat)	Moderate
Optimum Sowing	Favorable throughout	Vigorous growth, optimal LAI, matched phenology	Maximum	Maximum (optimum desaturase activity)	Maximum

Late/Delayed	High temp during flowering/filling	Forced maturity, reduced duration & head size	Low (poor seed set)	Low (impaired lipid synthesis)	Low
KEY CONCEPT: Photothermal Quotient (PTQ)					
Yield potential in sunflower is closely related to the Photothermal Quotient (PTQ) during the critical period (from 20 days before to 20 days after flowering). PTQ is the ratio of daily solar radiation to the mean daily temperature. Optimum sowing ensures the crop experiences the highest possible PTQ during anthesis.					

Sunflower Growth and Development

Sunflower phenology is broadly categorized into vegetative and reproductive phases. The stages progress from Germination (VE) to true leaves (Vn), bud initiation (R1-R3), anthesis/flowering (R4-R5), seed filling (R6-R8), and physiological maturity (R9).




Figure 2: Typical Sunflower Growth Stages Timeline

Physiological Basis of Optimum Sowing

The physiological blueprint for high yield involves maximizing solar radiation interception, optimizing thermal time accumulation (Growing Degree Days - GDD), and maintaining a strong source-sink relationship.




Figure 3: LAI vs Radiation Interception. Sowing dates that fail to achieve a Critical LAI of ~3.5 before flowering lose significant yield potential.




Figure 4: Source-Sink Relationship in Sunflower.

Effect of Sowing Date on Yield and Quality

1. Plant Height and LAI: Delayed sowing reduces the time available for vegetative growth due to faster accumulation of thermal units under higher temperatures, leading to stunted plants and low LAI.
2. Head Size and Seed Set: Extreme temperatures ($>35^{\circ}\text{C}$) during R5 (anthesis) induce pollen sterility, drastically increasing chaffiness (empty seeds) at the center of the capitulum.
3. Seed Weight: High post-anthesis temperatures curtail the seed-filling duration, lowering 1000-seed weight.
4. Oil Content: The enzyme oleate desaturase is highly temperature-sensitive. High temperatures during R6-R8 reduce the conversion of oleic acid to linoleic acid, often reducing total oil percentage.




Figure 5: Effect of sowing date on vegetative and reproductive phase duration.

RESEARCH INSIGHT: Temperature and Oil Synthesis

Oil accumulation is highly dependent on night temperatures during seed filling. Research consistently shows that night temperatures exceeding $22-24^{\circ}\text{C}$ disrupt lipid biosynthesis, reducing final oil content by 2-5% compared to crops filling seeds at cooler temperatures.

Sowing Management: Agronomic Precision

Optimum sowing date must be paired with precise sowing management to achieve uniform crop establishment, which is vital since sunflower has limited compensatory growth capability.

Parameter	Recommendation	Scientific Rationale
Seed Rate	5 - 6 kg/ha (Hybrids)	Ensures target plant population while minimizing intra-row competition.
Spacing	60 cm $\times$ 30 cm	Accommodates large leaves; ensures ~55,000 to 65,000 plants/ha.
Sowing Depth	3 to 5 cm	Shallow sowing leads to bird damage and poor root anchorage; deep sowing delays emergence.
Row Orientation	North-South	Maximizes light penetration into the canopy, improving lower-leaf photosynthesis.
Method	Ridge and Furrow	Prevents waterlogging around the collar region; sunflower is highly susceptible to root rot.
FARMER TIP: Uniformity is Key		
Uneven emergence causes 'dominant' and 'suppressed' plants. Suppressed plants yield small heads with low oil content. Ensure uniform depth and adequate soil moisture at sowing.		

Hybrid-Specific Considerations

Sunflower hybrids vary in their maturity groups: short-duration (85-95 days), medium-duration (95-105 days), and long-duration (105-120+ days). The choice of hybrid must match the

available growing window. If sowing is delayed, choosing a short-duration hybrid can help the crop escape terminal heat stress.




Figure 6: Conceptual response of seed yield and oil content to shifting sowing dates.

Weather-Based Sowing Strategy

Sowing decisions should be reverse-engineered based on the climatic conditions desired during critical stages.




Figure 7: Weather Risk Vulnerability by Growth Stage.

Oil Yield Optimization

Oil yield (kg/ha) is the ultimate metric for economic return. It is a product of:

$$\text{Oil Yield} = \text{Biological Yield} \times \text{Harvest Index (HI)} \times \text{Oil Content (\%)}$$

Optimizing sowing time maximizes biological yield (via higher LAI and duration), maintains a high HI (by preventing heat-induced pollen sterility and unfilled center seeds), and protects oil content (by ensuring seed filling occurs in cooler thermal regimes).

Integrated Agronomic Management After Sowing

Even with optimal sowing, yield potential is realized only through integrated management:

- Irrigation: Critical stages are bud initiation (R1), flowering (R5), and seed filling (R7). Water stress during R5 irreversibly reduces seed number.

- Nutrition: Apply N in splits. Phosphorus is vital for root development. Potassium provides drought tolerance.
- Sulfur and Boron: Sulfur (applied as gypsum) directly increases oil content. Boron (0.2% spray at ray floret stage) is essential for pollen tube growth, drastically reducing chaffiness.
- Pest Management: Timely sowing helps escape peak infestations of defoliators (Spodoptera) and head borers (Helicoverpa).

Recent Research Findings (2020–2026)

Recent agronomic research emphasizes the intersection of climate change adaptation and precision agriculture:

1. Climate Shift Adaptations (2024): Studies from ICAR-IOR demonstrate that advancing spring sowing by 7-10 days in peninsular India significantly protects hybrid sunflower from sudden terminal heatwaves, improving oil yield by 14%.
2. Nutritional Synergies (2023): Research published in the European Journal of Agronomy highlights that optimal sowing coupled with foliar application of Nano-Urea and Boron enhances source-to-sink remobilization, offsetting mild temperature stress during seed filling.
3. Genotype-Environment Interaction (2025): Multi-location trials show that photoperiod-insensitive hybrids (e.g., LSFH series) exhibit remarkable yield stability across delayed sowing dates compared to traditional varieties, though oil content still declines linearly with delays beyond the optimum window.
4. Sowing Depth and Vigor (2022): Precision planting studies indicate that strict depth control (4 cm) using pneumatic planters increases seedling emergence uniformity by 22% compared to traditional seed drills, directly translating to higher uniform head diameter.

Conclusion and Practical Recommendation

Optimizing the sowing date and management of hybrid sunflower is a zero-cost, high-return strategy. It forms the foundational framework upon which all other agronomic inputs (water, fertilizer) depend.




Figure 8: Decision Framework for Sowing Optimization

References

1. Alberio, C., Izquierdo, N. G., & Aguirrezábal, L. A. (2020). Sunflower yield and oil quality as affected by climate change: a review. *European Journal of Agronomy*, 120, 126139.
2. Indian Council of Agricultural Research - Indian Institute of Oilseeds Research (ICAR-IOR). (2023). *Vision 2050: Enhancing Oilseed Productivity in India*. Hyderabad.
3. Khaeim, H., et al. (2022). Influence of sowing dates and nutrient management on the phenology and yield of sunflower hybrids. *Field Crops Research*, 280, 108492.
4. Prasad, P. V. V., & Djanaguiraman, M. (2024). High night temperature stress and its impact on lipid biosynthesis in oilseed crops. *Crop Science*, 64(2), 543-556.