



Molecular Identification of Morphologically Similar Insect Species: Opportunities and Limitations

*Darshana Brahma and Pranita Roy

Ph.D. Scholar, ICAR-National Bureau of Agricultural Insect Resources,
Bengaluru, Karnataka, India

*Corresponding Author's email: darshanabrahma1@gmail.com

Correct identification of an insect is often the first step towards its successful management. In agricultural entomology, species identity influences decisions related to pest control, biological control, quarantine, invasive-pest surveillance and management of insect vectors. Misidentification can lead to inappropriate management practices, incorrect pest records and even mismatches between biological-control agents and their intended targets (Cock, 2015). Traditionally, insects are identified using morphological characters such as body structure, colour, wing venation, antennae, chaetotaxy, genitalia and features of immature stages. Morphological taxonomy remains fundamental to species recognition and classification; however, identification may become difficult when diagnostic characters are minute, specimens are immature or damaged, or closely related species are morphologically very similar (Schiff et al., 2012). The problem becomes particularly important in cryptic species, where biologically distinct species possess few obvious morphological differences. Such species may nevertheless differ in host preference, geographical distribution, insecticide susceptibility, invasive ability or capacity to transmit pathogens (Bickford et al., 2007).

A well-known agricultural example is the *Bemisia tabaci* species complex. Members of this complex are extremely difficult to distinguish morphologically, although molecular studies have revealed considerable genetic and biological differentiation among them (De Barro et al., 2011). Similar difficulties occur among economically important fruit flies, for which morphological, molecular, behavioural and ecological evidence has been combined to resolve *Anastrepha fraterculus* complex, *Ceratitis* FAR (*C. fasciventris*, *C. anonae*, *C. rosa*) complex and *Bactrocera dorsalis* complex (Schutze et al., 2017).

Molecular techniques therefore provide an important additional layer of evidence. DNA barcoding, PCR-based diagnostics and sequencing of mitochondrial and nuclear genes have become valuable tools for confirming species identity, detecting cryptic diversity and identifying specimens that cannot be reliably diagnosed by morphology alone.

DNA barcoding: A molecular identity tag

DNA barcoding uses a short, standardized DNA sequence to assist species identification. Hebert et al. (2003) proposed a portion of the mitochondrial cytochrome c oxidase subunit I (*COI*) gene as a standard barcode for animals. In a typical workflow, DNA is extracted from an insect, the target gene is amplified by PCR and the resulting sequence is compared with authenticated reference sequences. Databases such as the Barcode of Life Data System (BOLD) link barcode sequences with voucher specimens, taxonomy and collection information (Ratnasingham and Hebert, 2007). The approach is particularly useful for insects because very small quantities of tissue are often sufficient for obtaining DNA. For Indian entomology, Shashank et al. (2022) emphasized both the increasing usefulness of DNA barcoding and the need to expand well-curated reference libraries for the country's exceptionally rich insect fauna.

Molecular markers used in insect identification

Although *COI* is the most familiar molecular marker, it is not the only gene useful for insect identification.

1. **Mitochondrial markers:** *COI* is widely used for routine DNA barcoding because it often contains enough sequence variation to separate closely related species and has extensive representation in public databases.
2. **Nuclear ribosomal markers:** Genes and regions such as 28S rRNA, 18S rRNA, ITS1, and ITS2 provide independent nuclear evidence. They are particularly useful when mitochondrial results are uncertain (Caterino et al., 2000; Jenkins et al., 2012).
3. **Multiple loci:** Analysis of more than one independent gene can increase confidence in difficult species-delimitation problems. Dupuis et al. (2012) showed that additional loci generally improve the reliability of delimiting closely related species.

Table 1. Comparison of morphological and molecular approaches for insect identification

Aspect	Morphological identification	Molecular identification
What is examined?	Visible body characters	DNA sequence
Examples	Wings, antennae, colour, genitalia, chaetotaxy	<i>COI</i> , 28S rRNA, ITS, 18S rRNA
Best suited for	Well-described species with clear diagnostic characters	Cryptic, immature, damaged or closely related species
Immature stages	Often difficult to identify	Usually possible if good-quality DNA is obtained
Damaged specimens	Difficult if key structures are missing	Often possible if DNA is preserved
Cryptic species	May remain unresolved	Often gives better discrimination
Expertise needed	Trained taxonomist	Molecular laboratory and sequence-analysis skills
Main advantage	Simple, direct and cost-effective	High resolving power for difficult taxa
Main limitation	Similar-looking species may be confused	Depends on correct reference sequences and suitable markers
Most reliable use	Together with molecular evidence	Together with morphology and biological information

Where molecular identification helps agriculture

1. Cryptic pest species

Molecular methods can reveal genetically distinct pests concealed under a single morphological name. This is important because closely related species may respond differently to pesticides, natural enemies and host plants. DNA barcoding has, for example, been successfully applied to discriminate members of the *B. tabaci* complex (Ashfaq et al., 2014).

2. Immature and damaged specimens

Many insect keys depend on adult characters. Eggs, larvae, nymphs or broken specimens recovered from traps and consignments may therefore be difficult to identify. DNA-based methods can often provide identification from such material as long as adequate DNA remains.

3. Quarantine and biosecurity

International movement of agricultural commodities creates opportunities for exotic pests to enter new regions. Rapid identification of an intercepted insect may determine whether quarantine action is necessary. Armstrong and Ball (2005) highlighted DNA barcoding as a valuable tool for invasive-species identification in biosecurity. Similarly, Ashfaq and Hebert (2016) demonstrated the importance of reference DNA barcodes for regulated and economically important arthropod plant pests.

4. Identification of vectors

Correct species identification is particularly important for insects transmitting plant viruses, phytoplasmas and other pathogens. Morphologically similar whiteflies, aphids, thrips or leafhoppers may vary in their vector competence. Molecular confirmation therefore strengthens studies of disease epidemiology (Bertin and Bosco, 2013).

5. Biological control

Parasitoids and predators may themselves contain cryptic species. Molecular identification can help confirm the identity of natural enemies and associate parasitoids with their correct hosts, supporting safer and more effective biological-control programmes.

Limitations of molecular identification

Molecular identification is powerful, but it is not automatically correct.

1. Reference databases can contain errors

Sequence comparison is only as reliable as the reference sequence against which an unknown specimen is compared. If the original voucher specimen was incorrectly identified, the incorrect name may enter the database. Collins and Cruickshank (2013) therefore emphasized the need for accurately identified voucher material when constructing barcode libraries.

2. A single gene cannot resolve every species

Although *COI* performs well in many insects, there is no universal percentage of sequence difference that separates all species. Intraspecific and interspecific variation can overlap. Meier et al. (2006), using a large dataset of Diptera, showed that *COI*-based identification can become difficult where within-species variation is high or closely related species have very similar sequences. Mitochondrial introgression, hybridization and incomplete lineage sorting may also result in closely related species sharing mitochondrial sequences.

3. Molecular identification and species discovery are different

Matching an unknown specimen to a reliable reference sequence is species identification. Finding a divergent DNA sequence, however, does not automatically mean that a new species has been discovered. Species delimitation requires broader evidence. Fixed genetic-distance thresholds should not be applied universally for species identification, because levels of sequence divergence can vary among taxa depending on factors such as population size, mutation rate and time since speciation (Collins and Cruickshank, 2013).


Integrative taxonomy: Combining the evidence

For morphologically similar or cryptic insects, integrative taxonomy provides a more reliable approach by combining morphology with molecular, ecological and biological information. Using multiple sources of evidence reduces the limitations of relying on a single method and improves species delimitation (Schlick-Steiner et al., 2010; Sheth and Thaker, 2017).

A reliable diagnosis may therefore combine:

Morphology + DNA sequences + host association + geographical distribution + ecology + behaviour

Molecular techniques should consequently be viewed as partners of classical taxonomy rather than replacements for it.






Figure 1. Proposed integrative workflow for reliable identification of morphologically similar insect species. Molecular evidence is used to complement morphological diagnosis, particularly when conventional characters are insufficient.

Future prospects

The future of insect identification is moving beyond conventional sequencing. High-throughput sequencing, metabarcoding, genome-wide markers and portable molecular platforms are increasingly expanding diagnostic possibilities. Bulk trap catches may eventually be screened simultaneously for multiple pest species, while rapid species-specific assays can support surveillance at quarantine stations or closer to production areas. For countries with high insect diversity, including India, the immediate priority remains development of well-curated molecular reference libraries linked to properly identified voucher specimens.

Conclusion

Molecular identification has greatly improved our ability to recognize insects that are difficult to separate morphologically. DNA barcoding and related tools are particularly valuable for cryptic species, immature stages, damaged specimens, quarantine interceptions and insect vectors. However, molecular data are not free from error. Their reliability depends on the quality of reference specimens, appropriate marker selection and careful interpretation. For agricultural entomology, the strongest approach is therefore not morphology versus molecules, but morphology supported by molecules. Integrating conventional taxonomy with molecular, ecological and biological information provides a more dependable basis for pest identification, quarantine decisions and sustainable pest management.

References

1. Armstrong, K.F. and Ball, S.L. (2005). DNA barcodes for biosecurity: invasive species identification. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **360**(1462): 1813-1823.
2. Ashfaq, M., Hebert, P.D.N., Mirza, M.S., Khan, A.M., Mansoor, S., Shah, G.S. and Zafar, Y. (2014). DNA barcoding of *Bemisia tabaci* complex (Hemiptera: Aleyrodidae) reveals southerly expansion of the dominant whitefly species on cotton in Pakistan. *PLoS ONE*, **9**(8): e104485.
3. Ashfaq, M. and Hebert, P.D.N. (2016). DNA barcodes for bio-surveillance: regulated and economically important arthropod plant pests. *Genome*, **59**(11): 933-945.
4. Bertin, S. and Bosco, D. (2013). Molecular identification of phytoplasma vector species. *Methods in Molecular Biology*, **938**: 87-108.
5. Bickford, D., Lohman, D.J., Sodhi, N.S., Ng, P.K.L., Meier, R., Winker, K., Ingram, K.K. and Das, I. (2007). Cryptic species as a window on diversity and conservation. *Trends in Ecology & Evolution*, **22**(3): 148-155.
6. Caterino, M.S., Cho, S. and Sperling, F.A.H. (2000). The current state of insect molecular systematics: a thriving tower of Babel. *Annual Review of Entomology*, **45**: 1-54.

7. Cock, M.J.W. (2015). A critical review of the literature on the pest *Erionota* spp. (Lepidoptera, Hesperidae): taxonomy, distribution, food plants, early stages, natural enemies and biological control. *CAB Reviews*, **10**(007): 1-30.
8. Collins, R.A. and Cruickshank, R.H. (2013). The seven deadly sins of DNA barcoding. *Molecular Ecology Resources*, **13**(6): 969-975.
9. De Barro, P.J., Liu, S.S., Boykin, L.M. and Dinsdale, A.B. (2011). *Bemisia tabaci*: a statement of species status. *Annual Review of Entomology*, **56**: 1-19.
10. Dupuis, J.R., Roe, A.D. and Sperling, F.A.H. (2012). Multi-locus species delimitation in closely related animals and fungi: one marker is not enough. *Molecular Ecology*, **21**(18): 4422-4436.
11. Hebert, P.D.N., Cywinska, A., Ball, S.L. and deWaard, J.R. (2003). Biological identifications through DNA barcodes. *Proceedings of the Royal Society B: Biological Sciences*, **270**(1512): 313-321.
12. Jenkins, C., Chapman, T.A., Micallef, J.L. and Reynolds, O.L. (2012). Molecular techniques for the detection and differentiation of host and parasitoid species and the implications for fruit fly management. *Insects*, **3**(3): 763-788.
13. Meier, R., Shiyang, K., Vaidya, G. and Ng, P.K.L. (2006). DNA barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. *Systematic Biology*, **55**(5): 715-728.
14. Ratnasingham, S. and Hebert, P.D.N. (2007). BOLD: The Barcode of Life Data System. *Molecular Ecology Notes*, **7**(3): 355-364.
15. Sheth, B. P. and Thaker, V. S. (2017). DNA barcoding and traditional taxonomy: an integrated approach for biodiversity conservation. *Genome*, **60**(7), 618-628.
16. Schiff, N.M., Goulet, H., Smith, D.R., Boudreault, C., Wilson, A.D. and Scheffler, B.E. (2012). Siricidae (Hymenoptera: Symphyta: Siricoidea) of the Western Hemisphere. *Canadian Journal of Arthropod Identification*, **21**: 1-305.
17. Schlick-Steiner, B.C., Steiner, F.M., Seifert, B., Stauffer, C., Christian, E. and Crozier, R.H. (2010). Integrative taxonomy: a multisource approach to exploring biodiversity. *Annual Review of Entomology*, **55**: 421-438.
18. Schutze, M.K., Virgilio, M., Norrbom, A. and Clarke, A.R. (2017). Tephritid integrative taxonomy: where we are now, with a focus on the resolution of three tropical fruit fly species complexes. *Annual Review of Entomology*, **62**: 147-164.
19. Shashank, P.R., Naveena, N.L., Rajgopal, N.N., Elliott, T.A., Sreedevi, K., Sunil, S. and Meshram, N.M. (2022). DNA barcoding of insects from India: current status and future perspectives. *Molecular Biology Reports*, **49**(11): 10617-10626.