



Beyond Planting Trees: The New Science of Mangrove Restoration

Sneha R Chaudhari, *Bindiya Parmar, Vanshita Tandel and D T Vaghela

College of Fisheries Science, Kamdhenu University, Veraval-362265, India

*Corresponding Author's email: bkparmar1330@gmail.com

Mangrove restoration is no longer just about planting more trees; it is about creating the right conditions for mangroves to grow, recover, and sustain themselves naturally. This article highlights the importance of healthy tidal flow, suitable hydrology and sediments, proper species selection, natural regeneration, biodiversity, and community involvement. Mangroves also provide valuable benefits by storing blue carbon, protecting coastlines, and supporting fisheries. New technologies such as GIS, remote sensing, drones, LiDAR, and AI are helping researchers monitor restoration more effectively. However, poor site selection, disrupted tidal flow, coastal development, and limited long-term monitoring can reduce restoration success. In the long term, restoration efforts should focus on healthy ecosystems, biodiversity, carbon sequestration, and lasting sustainability rather than simply counting planted seedlings.

Keywords: Blue carbon; Carbon sequestration; Ecosystem sustainability; Mangrove; Restoration

“Mangrove restoration is not just about planting trees; it is about restoring nature’s ability to protect coasts, support biodiversity and store carbon for generations.”

Introduction

Mangroves are special coastal ecosystems that play an important role in protecting shorelines, supporting biodiversity and sustaining fisheries. They also store large amounts of carbon, particularly in their sediments, making them important ecosystems for climate-change mitigation. In addition, mangrove areas provide nursery and feeding grounds for many fish and invertebrates and support the livelihoods and food security of coastal communities (Donato et al., 2011). However, mangroves are facing increasing pressure from coastal development, aquaculture, pollution and changes in natural water flow. In the past, restoration efforts mainly focused on planting large numbers of seedlings. Although planting can help in suitable locations, it may not succeed when tidal flow, elevation, sediment conditions or other environmental factors are inappropriate. Monoculture plantations may also fail to reproduce the diversity and resilience of natural mangrove forests (Lewis & Brown, 2014). For this reason, mangrove restoration is gradually moving beyond simply planting trees. Modern approaches focus on restoring the natural conditions that allow mangroves to regenerate and grow on their own.

Mangrove Planting to Ecosystem Restoration

Mangrove restoration is no longer seen as simply planting as many seedlings as possible. The main aim is to restore the natural conditions that help mangroves grow, spread, and regenerate naturally (Lewis & Brown, 2014). Natural recovery depends on healthy tidal flow, suitable soil elevation, salinity, and sediment conditions. Roads, embankments, and aquaculture activities can disturb these processes. Restoring natural water movement can therefore help mangroves recover more effectively (Beeston et al., 2023).

The Science Behind Successful Mangrove Restoration

Mangrove restoration is not just about planting more trees along the coast. It means restoring the natural conditions that help mangroves grow, regenerate, and survive in the long term. Healthy tidal flow, suitable sediments, native mangrove species, natural regeneration, biodiversity, and support from local communities are all important for successful restoration (Su et al., 2021). Tides are especially important because they bring water, nutrients, and sediments into mangrove habitats. When roads or other structures block these natural water pathways, mangrove growth can be disturbed. Restoring tidal connections can therefore help mangroves recover naturally (Lewis et al., 2018). Similarly, stable sediments and the use of native species adapted to local salinity and flooding conditions can improve mangrove establishment (Ray et al., 2024). When environmental conditions are suitable, allowing mangroves to regenerate naturally may be more effective than planting many seedlings (Su et al., 2021). The return of fish, crabs, birds, and other organisms can also show that the ecosystem is recovering (Lovelock et al., 2025). Local communities can contribute valuable knowledge, making natural ecosystem recovery more important than tree planting alone.

Mangroves and Blue Carbon: Beyond Tree Planting

Understanding Blue Carbon: Blue carbon is the carbon captured and stored by coastal ecosystems such as mangroves, tidal marshes, and seagrass meadows. Mangroves are especially valuable because they store large quantities of carbon not only in their trees and roots but also in the waterlogged soils beneath them. In contrast to many terrestrial forests, where much of the carbon is stored in aboveground vegetation, a major share of mangrove carbon is found below the ground in roots and sediments (Macreadie et al., 2021).




Figure 1. Blue carbon flux and soil accumulation. Source: Adapted from the Manual for the Creation of Blue Carbon Projects in Europe and the Mediterranean, LIFE Blue Natura Project (LIFE14 CCM/ES/000957).

Carbon Storage and Long-Term Burial: Carbon in mangrove ecosystems is stored in aboveground vegetation, belowground roots, and coastal sediments. Among these pools, the sediment is particularly important because regular tidal flooding keeps the soil wet and often limits the availability of oxygen. This slows the decomposition of organic matter and allows carbon from fallen leaves, roots, and other organic materials to accumulate in the soil. When these sediments remain undisturbed, carbon can stay buried for long periods. However, activities such as clearing, drainage, erosion, or other forms of disturbance can expose stored carbon and increase greenhouse-gas emissions (Adame et al., 2021).

Restoration, Recovery and Carbon Sequestration: Restoring mangroves can help rebuild carbon stocks, but restoration is not simply a matter of planting large numbers of seedlings. In the early stages of recovery, carbon accumulation mainly occurs as vegetation and root systems develop. As the mangrove grows, roots help stabilise sediments, and organic material gradually builds up in the soil. Over time, these processes contribute to the development of a larger belowground carbon pool. However, restored mangroves may need several decades to approach the carbon stocks and ecological functions of mature mangrove forests (Su et al., 2021). Therefore, restoration should focus on restoring suitable hydrology, encouraging natural regeneration, selecting appropriate species, and matching restoration activities to local site conditions.

Protection and Restoration: Protecting existing mangrove forests is just as important as restoring degraded areas. Mature mangroves already contain substantial amounts of carbon accumulated over many years. When these ecosystems are cleared or disturbed, both vegetation and carbon-rich sediments can be lost, potentially releasing stored carbon back into the

atmosphere. Restoration can help recover degraded ecosystems, but preventing the destruction of healthy mangroves avoids the loss of carbon stocks that may take many years or decades to rebuild (Macreadie et al., 2021).

Mangrove Carbon: Monitoring mangrove blue carbon requires a combination of field studies, GIS, and remote-sensing tools. Soil samples can be used to estimate carbon stored in sediments, while tree measurements help assess above-ground biomass. Technologies such as UAV LiDAR, photogrammetry, Sentinel-2, and Landsat make it possible to study mangrove cover and changes over larger areas (Su et al., 2021). Today, mangrove restoration is moving beyond simply planting trees. The focus is on restoring natural conditions that help mangroves grow, store carbon, support wildlife, and protect coastal areas. This approach can contribute to healthier and more resilient coastal ecosystems.

Technology-Based Monitoring of Mangrove Restoration

Mangrove restoration is no longer judged only by the number of seedlings planted. A site may have many surviving plants but still fail to develop into a healthy and self-sustaining ecosystem. Modern restoration therefore focuses on long-term changes in vegetation, hydrology, biodiversity, ecosystem structure, and carbon storage. Technologies such as GIS, satellite remote sensing, drones, LiDAR, and artificial intelligence (AI) are making this monitoring faster and more reliable (Yang et al., 2024).

GIS and Satellite Remote Sensing: GIS and satellite imagery allow large mangrove areas to be monitored repeatedly over time. Landsat and Sentinel-2 images can show changes in mangrove cover, vegetation condition, fragmentation, and recovery. Sentinel-1 radar data are especially useful in cloudy coastal regions and can provide additional information on vegetation structure and surface conditions. GIS can also combine elevation, tidal channels, shoreline change, and land-use information to identify suitable restoration sites. This is important because successful mangrove recovery depends strongly on appropriate tidal and hydrological conditions, not simply on planting more trees (Yang et al., 2024).

Drone-Based Monitoring: Drones provide much finer information than satellites and are useful for monitoring small restoration sites. High-resolution RGB, multispectral, and LiDAR images can help measure seedling survival, canopy development, tree height, and gaps within the vegetation. UAV-based LiDAR can also provide detailed information about vegetation structure and help estimate aboveground biomass and carbon (Duan et al., 2025). Field observations are still important, but drones can greatly reduce the time and effort required for repeated surveys.

Vegetation and Habitat Mapping: Restoration success also depends on whether the restored area develops a diverse and natural vegetation structure. Multispectral and hyperspectral images can help distinguish mangroves from other vegetation and, in some cases, identify different mangrove species. Vegetation indices such as NDVI and EVI can be used to monitor changes in canopy condition and plant growth. Recent developments in remote sensing and machine learning have improved the mapping of mangrove species and vegetation patterns (Wu et al., 2025).

AI and Machine Learning: Large amounts of satellite and drone data can be difficult to analyse manually. Machine-learning methods such as Random Forest, Support Vector Machine, and deep-learning models can automatically classify mangroves, detect changes, and estimate biomass. These approaches can also combine optical, radar, elevation, and LiDAR data to provide a more detailed picture of ecosystem recovery (Wu et al., 2025). However, AI-based results should always be checked with field observations to ensure reliable interpretation.

Biodiversity and Blue Carbon: Modern restoration monitoring is increasingly concerned with ecosystem functions, not just plant survival. Changes in species diversity, natural regeneration, habitat quality, and associated fish, birds, and invertebrates can indicate whether ecological recovery is taking place. Blue carbon is another important indicator because mangroves store carbon in vegetation, roots, and coastal sediments. Remote sensing and LiDAR can help estimate aboveground biomass, while field and sediment measurements are needed to assess belowground and soil carbon (Duan et al., 2025).

Long-Term Monitoring and Adaptive Management: The real success of restoration can only be understood through long-term monitoring. Repeated observations can show whether mangroves are developing greater canopy cover, natural regeneration, structural complexity, biodiversity, and carbon-storage capacity. They can also reveal problems such as erosion, poor tidal connectivity, or vegetation decline at an early stage. This information allows managers to modify restoration practices when necessary. Overall, the combination of field surveys, GIS, satellite imagery, drones, LiDAR, and AI is helping move mangrove restoration beyond simple tree planting. The goal is not only to increase mangrove cover, but to develop healthy, diverse, functional, and carbon-rich ecosystems that can persist over the long term (Yang et al., 2024).

Challenges and Future Directions

Mangrove restoration can be difficult when the wrong species are selected, tidal flow is disturbed, drainage is poor, or coastal habitats are lost to development (Su et al., 2021). Another major issue is long-term monitoring. Simply checking how many planted seedlings survive is not enough to understand whether the ecosystem is recovering. Changes in biodiversity, natural regeneration, carbon storage, and habitat quality should also be followed over time (Sasmito et al., 2023). In the future, restoration should focus less on the number of trees planted and more on how well the ecosystem functions. Protecting biodiversity, storing carbon, stabilising shorelines and sediments, and supporting fisheries can provide better measures of successful restoration (Lovelock et al., 2025).

Conclusion

Mangrove restoration is not simply about planting more trees; it is about creating the right conditions for mangroves to grow, regenerate, and function as healthy coastal ecosystems. When natural tidal flow, suitable habitats, biodiversity, and sediment processes are restored, mangroves can provide long-term benefits such as coastal protection, carbon storage, and habitat for many species (Su et al., 2021). In the future, restoration should focus on healthy and self-sustaining ecosystems rather than plantation numbers alone. Giving importance to natural regeneration, biodiversity, carbon sequestration, community involvement, and long-term monitoring can make restoration efforts more effective and lasting. In this way, mangrove restoration can contribute not only to coastal conservation but also to the well-being and resilience of coastal communities (Lovelock et al., 2025).

References

1. Adame, M. F., Connolly, R. M., Turschwell, M. P., Lovelock, C. E., Fatoyinbo, T., Lagomasino, D., Goldberg, L. A., Holdorf, J., Friess, D. A., Sasmito, S. D., Sanderman, J., Sievers, M., Buelow, C., Simard, M., & Brown, C. J. (2021). Future carbon emissions from global mangrove forest loss. *Global Change Biology*, 27(12), 2856–2866. <https://doi.org/10.1111/gcb.15571>
2. Alongi, D. M. (2020). Global significance of mangrove blue carbon in climate change mitigation. *Scientific Reports*, 10, 10057. <https://doi.org/10.1038/s41598-020-71103-0>
3. Beeston, M., Cameron, C., Hagger, V., Howard, J., Lovelock, C., Sippon, J., Tonneijck, F., van Bijsterveldt, C., & van Eijk, P. (Eds.). (2023). *Best practice guidelines for mangrove restoration*. Global Mangrove Alliance. Official restoration guidelines
4. Budiarto, R., & Dewanto, B. G. (2025). Unmanned aerial vehicles for assessing biomass and carbon stocks in mangrove forests: A systematic review. *Sustainable Futures*, 10, 101425. <https://doi.org/10.1016/j.sfr.2025.101425> ScienceDirect
5. Castillo, J. A. A., et al. (2022). A systematic review of mangrove restoration studies in Southeast Asia: Challenges and opportunities for the United Nations Decade on Ecosystem Restoration. *Frontiers in Marine Science*, 9, 987737. Frontiers
6. Charoenlerkthawin, W., Bidorn, K., Burnett, W. C., Sopon, A., Otarawanna, S., & Bidorn, B. (2024). Evaluating the effectiveness of mangrove rehabilitation: A novel approach for sustainable coastal management. *Journal of Environmental Management*, 365, 121673. <https://doi.org/10.1016/j.jenvman.2024.121673> PubMed

7. Donato, D. C., Kauffman, J. B., Murdiyarso, D., Kurnianto, S., Stidham, M., & Kanninen, M. (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*, 4(5), 293–297. <https://doi.org/10.1038/ngeo1123>
8. Duan, M., Sánchez-Azofeifa, A., Abdulmajeed, M., Turner, D. P., Buckingham, K., Odari, A., Mtwana, J., Kipkoech, S., & Kasraee, N. (2025). Aboveground carbon estimation in a mangrove ecosystem using UAV-based remote sensing and machine learning. *Ecological Indicators*, 178, 113950. <https://doi.org/10.1016/j.ecolind.2025.113950> ScienceDirect
9. Friess, D. A., Yando, E. S., Alemu I, J. B., Wong, L.-W., Bhatia, N., & Yaakub, S. M. (2020). Mangrove rehabilitation and restoration as experimental adaptive management. *Frontiers in Marine Science*, 7, 327. Frontiers
10. Global Mangrove Alliance & Blue Carbon Initiative. (2023). *Best practice guidelines for mangrove restoration*. Global Mangrove Alliance.
11. Lewis, R. R. (2005). Ecological engineering for successful management and restoration of mangrove forests. *Ecological Engineering*, 24(4), 403–418. <https://doi.org/10.1016/j.ecoleng.2004.10.003> reefresilience.org
12. Lewis, R. R., Brown, B. M., & Flynn, L. L. (2018). Methods and criteria for successful mangrove forest rehabilitation. In *Coastal wetlands: An integrated ecosystem approach* (2nd ed.). Elsevier. SDG Resource Centre
13. Lewis, R. R., III, & Brown, B. (2014). *Ecological mangrove rehabilitation: A field manual for practitioners*. Mangrove Action Project. FAO resource page
14. Lovelock, C. E., Hagger, V., Feller, I. C., Amir, A. A., António, V. M., Owuor, M. A., & Friess, D. A. (2025). Mangrove biodiversity and ecosystem services. *Nature Reviews Biodiversity*, 1, 772–787. <https://doi.org/10.1038/s44358-025-00103-3>
15. Macreadie, P. I., Costa, M. D. P., Atwood, T. B., Friess, D. A., Kelleway, J. J., Kennedy, H., Lovelock, C. E., Serrano, O., & Duarte, C. M. (2021). Blue carbon as a natural climate solution. *Nature Reviews Earth & Environment*, 2, 826–839. <https://doi.org/10.1038/s43017-021-00224-1>
16. Nascimento Neres, J., Strenzel, G. M. R., Mielke, M. S., & Barros, F. (2024). Mangrove forest health condition from space and the use of in situ data. *Marine Environmental Research*, 201, 106704. <https://doi.org/10.1016/j.marenvres.2024.106704> ScienceDirect
17. Pradisty, N. A., Schlund, M., Horstman, E. M., & Willemen, L. (2025). Estimating canopy height and aboveground biomass in tropical mangrove restoration areas through multisource remote sensing. *Ecological Informatics*, 92, 103522. <https://doi.org/10.1016/j.ecoinf.2025.103522> DOI
18. Ray, K., Basak, S. K., Giri, C. K., Kotal, H. N., Mandal, A., Chatterjee, K., Saha, S., Biswas, B., Mondal, S., Das, I., Ghosh, A., Bhadury, P., & Joshi, R. (2024). Ecological restoration at pilot-scale employing site-specific rationales for small-patch degraded mangroves in Indian Sundarbans. *Scientific Reports*, 14, 12952. <https://doi.org/10.1038/s41598-024-63281-8> Nature
19. Ray, K., et al. (2024). Ecological restoration at pilot-scale employing site-specific rationales for small-patch degraded mangroves in Indian Sundarbans. *Scientific Reports*, 14, 12952. <https://doi.org/10.1038/s41598-024-63281-8> Nature
20. Sasmito, S. D., Basyuni, M., Kridalaksana, A., Saragi-Sasmito, M. F., Lovelock, C. E., & Murdiyarso, D. (2023). Challenges and opportunities for achieving Sustainable Development Goals through restoration of Indonesia's mangroves. *Nature Ecology & Evolution*, 7, 62–70. Nature
21. Su, J., Friess, D. A., & Gasparatos, A. (2021). A meta-analysis of the ecological and economic outcomes of mangrove restoration. *Nature Communications*, 12, 5050. <https://doi.org/10.1038/s41467-021-25349-1>
22. Worthington, T. A., Andradi-Brown, D. A., Bhargava, R., Bunting, P., Duncan, C., Fatoyinbo, T., Lagomasino, D., Lovelock, C. E., Murray, N. J., Narayan, S., & Spalding, M. (2020). Harnessing big data to support the conservation and rehabilitation of mangrove forests globally. *One Earth*, 2(5), 429–443. <https://doi.org/10.1016/j.oneear.2020.04.018>

23. Wu, Y., Lu, C., Wu, K., Gao, W., Yang, N., & Lin, J. (2025). Advancements and trends in mangrove species mapping based on remote sensing: A comprehensive review and knowledge visualization. *Global Ecology and Conservation*, 57, e03408. <https://doi.org/10.1016/j.gecco.2025.e03408> ScienceDirect
24. Yang, X., Zhu, Z., Kroeger, K. D., Qiu, S., Covington, S., Conrad, J. R., & Zhu, Z. (2024). Tracking mangrove condition changes using dense Landsat time series. *Remote Sensing of Environment*, 315, 114461. <https://doi.org/10.1016/j.rse.2024.114461>