



Microbial Succession and Transformation of Organic Matter

^{*}Anisa N. A.¹ and Dr. Anu Rajan S.²

¹Assistant Professor, Agricultural Information and Sales Centre, Vengeri, Kozhikode, Kerala Agricultural University, India-673010

²Assistant Professor, Department of Agricultural Microbiology, College of Agriculture, Vellayani, Thiruvananthapuram, Kerala Agricultural University, India- 695522

^{*}Corresponding Author's email: anisa.na@kau.in

Soil is a living ecosystem where microorganisms act as nature's recyclers, transforming organic residues into simpler compounds and nutrients essential for plant growth. Microbial succession, the sequential change in microbial communities during decomposition, plays a crucial role in this transformation. This article explores the stages of microbial succession, the enzymes and microorganisms involved in organic matter decomposition, the processes governing nutrient cycling and carbon transformation, and the factors influencing these activities. Understanding these processes is essential for improving soil fertility, managing agricultural residues and promoting sustainable agriculture.

Soil organic matter: The foundation of a living soil

Soil organic matter is made up of both living and non-living components. The living fraction includes bacteria, fungi and other soil organisms, whereas the non-living fraction includes dead plant and animal materials, organic residues and excreta of soil organisms. Plant residues are particularly important because they contain several major organic compounds. Cellulose, hemicellulose and lignin form important structural components of plant tissues along with proteins, lipids, organic acids and phenolic compounds. The chemical composition of a residue strongly influences how quickly it decomposes. Simple compounds such as sugars and amino acids are readily utilized by microorganisms, whereas compounds like cellulose, lignin and other complex compounds require specialized enzymes and microorganisms for their breakdown. This explains why a freshly fallen leaf may disappear relatively quickly, while woody material can remain in the soil for much longer. The quality of organic material therefore matters just as much as the quantity added to the soil. Organic matter serves as a major reservoir of carbon, hydrogen and oxygen. Small amounts of nutrients such as nitrogen, phosphorous, sulphur, potassium, calcium and magnesium are also contained within organic residues. They are the primary source of carbon and nitrogen for the microbial communities. It improves soil structure, enhances water holding capacity and aeration, increases cation exchange capacity, supports an enormous diversity of soil microorganisms, improves water and air infiltration rate and plays a vital role in carbon sequestration, thereby contributing to climate change mitigation.

Decomposition: Nature's recycling system

When plant and animal residues enter the soil, microorganisms begin to colonize them. Decomposition involves the physical breakdown and biochemical transformation of complex organic molecules into simpler compounds. The process releases carbon dioxide, water, energy and nutrients. Under aerobic conditions, carbon is mainly released as carbon dioxide, whereas oxygen limited environments can promote anaerobic pathways that produce compounds such as methane and organic acids. Microorganisms decompose organic matter primarily to obtain energy for growth, carbon for cell formation and nutrients required for microbial growth.

Modern soil research increasingly recognizes that microorganisms do not merely destroy soil organic matter. Their growth, residues and transformation products can also contribute to the formation and stabilization of persistent soil organic matter. Therefore, the microbial community plays a dual role: it drives organic matter turnover while also contributing to long term carbon storage.

Microbial succession

Organic matter added to a field undergoes a continuous sequence of biological transformations. Microorganisms act as nature's recyclers, converting complex materials into simpler compounds and ultimately releasing nutrients that plants can use. But decomposition is not performed by a single group of microorganisms. Different microbial communities become important at different stages. This orderly change in microbial dominance is known as microbial succession. It is a dynamic ecological process which refers to the sequential replacement of microbial communities during decomposition as environmental conditions and substrate composition change. It ensures efficient utilization of all organic substrates and complete decomposition. One of the most fascinating features of decomposition is that microorganisms do not all become active at the same time. During the early stage of decomposition, microorganisms that can rapidly utilize simple compounds become dominant. As these compounds are exhausted, microorganisms capable of degrading more complex materials become increasingly important. Thus, microbial communities undergo a sequential replacement, or microbial succession, as the organic substrate and environmental conditions change. Microbial succession can broadly be viewed through four stages (Fig. 1).

Stage 1: Initial colonization

The first stage begins soon after fresh organic material enters the soil. At this stage, simple compounds such as sugars, amino acids, organic acids and soluble proteins are readily available. Fast growing bacteria and yeasts rapidly colonize the material. Genera such as *Bacillus*, *Pseudomonas* and *Enterobacter* can participate in this early phase. Microbial growth and respiration are high because easily available substrates provide an immediate source of energy. Nutrients are rapidly taken up by microorganisms, initiating the decomposition process. For agricultural residues rich in easily degradable compounds, this stage can proceed rapidly.




Fig. 1 Microbial succession of organic matter decomposition

Stage 2: The active decomposition phase

Once easily available compounds are depleted, microorganisms turn their attention towards more complex plant materials. Cellulose, hemicellulose and pectin become important substrates. This stage involves strong activity of cellulolytic bacteria and fungi. Microorganisms such as *Cellulomonas*, *Bacillus*, *Aspergillus*, *Penicillium* and *Trichoderma* are associated with the degradation of plant residues. This is often the most active phase of

decomposition. Microbial biomass increases and organic matter is rapidly transformed. The agricultural importance of this stage is considerable because nutrients contained within organic residues begin to move through the soil nutrient cycle.

Stage 3: The late decomposition

After easily degradable materials have been consumed, resistant compounds remain. Among the most important are lignin, waxes and polyphenolic compounds. These materials are chemically complex and cannot be easily degraded by all microorganisms. Specialized fungi, particularly lignin degrading fungi, become increasingly important. White rot fungi are particularly well known for their ability to attack lignin. Actinomycetes, including *Streptomyces*, also contribute to the degradation of resistant organic materials. At this stage, decomposition becomes slower. However, the process remains essential because it allows the remaining complex materials to undergo further transformation.

Stage 4: Humification

The final stage involves the transformation of resistant organic materials into more stable forms of soil organic matter. This process is known as humification. Actinomycetes and slow growing fungi participate in the transformation of remaining organic materials. The resulting humic substances contribute to the long term storage of carbon in soil. Humus improves several important soil properties, like water retention, nutrient retention, soil structure, cation exchange capacity and carbon storage. Thus, the end of decomposition is not simply the disappearance of organic matter. Some of the transformed material becomes part of the more stable organic matter pool.

Enzymes in organic matter decomposition

Microorganisms cannot simply absorb large molecules such as cellulose, proteins and lignin directly through their cell membranes. They therefore rely heavily on enzymes (Table 1). Some enzymes are constitutive enzymes meaning they are produced continuously by microbial cells regardless of the substrate present. In contrast, inducible enzymes are synthesized only when a specific substrate is available in the environment. Extracellular enzymes are secreted outside microbial cells and break large molecules into smaller compounds. These smaller molecules can then enter microbial cells and be metabolized. This makes enzymes the biochemical tools that drive decomposition. The production of different enzymes allows different microorganisms to specialize in different stages of decomposition.

Table 1. Different enzymes acting during decomposition and their main function

Enzyme	Function	Major microorganisms producing the enzymes
Cellulase	Cellulose degradation	<i>Trichoderma, Aspergillus, Penicillium, Cellulomonas, Bacillus, Clostridium</i>
Xylanase	Hemicellulose degradation	<i>Aspergillus, Trichoderma, Penicillium, Bacillus, Streptomyces</i>
Pectinase	Pectin degradation	<i>Aspergillus, Penicillium, Trichoderma, Bacillus, Pseudomonas</i>
Protease	Protein degradation	<i>Bacillus, Pseudomonas, Streptomyces, Aspergillus, Penicillium</i>
Lipase	Lipid degradation	<i>Pseudomonas, Bacillus, Aspergillus, Penicillium, Candida</i>
Laccase	Lignin oxidation	<i>Trametes, Pleurotus, Phanerochaete, Aspergillus, Penicillium</i>
Lignin peroxidase (LiP)	Lignin degradation	<i>Phanerochaete, Trametes, Pleurotus</i>
Manganese peroxidase (MnP)	Lignin degradation	<i>Phanerochaete, Trametes, Pleurotus</i>

Decomposition of major components of organic matter

Cellulose

Cellulose is one of the most abundant organic compounds in plant residues. It is composed of glucose units linked together in long chains. Microorganisms produce cellulase and related enzymes to break these chains into smaller molecules. The resulting sugars can then be absorbed and used for microbial energy and growth. Important cellulose degrading microorganisms include *Trichoderma*, *Cellulomonas*, *Phanerochaete* and *Clostridium* and several other bacteria and fungi. However, cellulose associated closely with lignin can be considerably more resistant to decomposition.

Hemicellulose

Hemicellulose is another important structural component of plant cell walls. Unlike cellulose, it has a more complex and branched structure. Its decomposition involves several hemicellulolytic enzymes. Microorganisms convert hemicellulose into simpler sugars such as xylose, arabinose, galactose and mannose. These compounds are subsequently metabolized by microorganisms. Fungi such as *Aspergillus*, *Penicillium*, *Trichoderma* and *Chaetomium*, together with bacteria such as *Bacillus* and *Pseudomonas*, can participate in hemicellulose decomposition.

Lignin

Lignin is one of the most resistant components of plant residues. Its complex aromatic structure makes it difficult for most microorganisms to degrade. Certain fungi possess powerful oxidative enzyme systems that enable them to attack lignin. Laccase, lignin peroxidase and manganese peroxidase are particularly important enzymes associated with lignin transformation. This is why materials containing high amounts of lignin generally decompose more slowly than materials containing large quantities of easily degradable compounds.

Proteins

Proteins contain nitrogen as well as carbon, hydrogen and oxygen. Their decomposition is therefore closely connected with the soil nitrogen cycle. Microorganisms such as *Bacillus*, *Pseudomonas*, *Aspergillus* and *Penicillium* produce proteases that break proteins into smaller compounds. The conversion of proteins into amino acids is called aminization, while the subsequent conversion of amino acids into ammonia is known as ammonification. The released nitrogen can then enter other stages of the nitrogen cycle and eventually become available to plants.

Chitin and lipids

Chitin occurs in fungal cell walls and the exoskeletons of arthropods. Although resistant to decomposition, it can be degraded by specialized microorganisms using enzymes such as chitinases. *Streptomyces* and *Nocardia*, along with fungi such as *Trichoderma*, participate in chitin decomposition. Lipids are another component of soil organic matter. Lipolytic microorganisms such as *Pseudomonas*, *Bacillus*, *Aspergillus* and *Penicillium* produce lipases and related enzymes that break down fats, oils and phospholipids. These processes return carbon and nutrients to the soil microbial system.

From decomposition to nutrient availability

The decomposition of organic matter is not simply the breakdown of dead plant and animal residues; it is a dynamic process through which microorganisms transform complex organic compounds into simpler forms and regulate the availability of nutrients in soil. The products of this decomposition may either be released into the soil as inorganic nutrients, temporarily incorporated into microbial biomass or transformed into more stable organic compounds. Thus, decomposition acts as a bridge between organic matter and plant available nutrients. Decomposition and transformation are closely connected with nutrient cycling.

Mineralization

Mineralization is the process by which microorganisms convert organically bound nutrients into inorganic forms that can be absorbed by plants. When microbes decompose nitrogen containing compounds such as proteins, amino acids and nucleic acids, organic nitrogen is

released as ammonium (NH_4^+). Under suitable conditions, nitrifying microorganisms subsequently convert ammonium into nitrite (NO_2^-) and then nitrate (NO_3^-), which are major forms of nitrogen available to plants. Similarly, decomposition releases phosphorus as inorganic phosphate and sulphur mainly as sulphate. Therefore, mineralization is an important microbial pathway through which nutrients locked in organic residues become available for plant uptake.

Immobilization

Immobilization temporarily reduces the availability of inorganic nutrients in soil. When microorganisms decompose residues with a high C:N ratio, such as straw, sawdust or mature crop residues, they require additional nitrogen to build their cells and reproduce. They therefore take up available inorganic nitrogen from the soil and incorporate it into their microbial biomass. This process is known as immobilization. Although immobilization may temporarily compete with plants for nutrients, it is not necessarily a loss. As microbial cells die and are further decomposed, the nutrients contained in their biomass can be released again through secondary mineralization. Immobilization therefore acts as a temporary nutrient storage mechanism and helps regulate nutrient cycling in soil. Thus, mineralization and immobilization work together to regulate nutrient availability in soil.

Humification

Humification represents another important pathway during decomposition. Not all organic matter is completely mineralized. Some partially decomposed plant residues, microbial products and complex organic compounds undergo further transformation, and contribute to the formation of humus and other stable forms of soil organic matter. Microbial activity breaks down easily degradable compounds first, while more resistant components, particularly lignin derived materials, persist longer and undergo chemical and biochemical transformations. The resulting humified organic matter improves soil structure, water holding capacity, cation exchange capacity and nutrient retention ability. Humus also provides a long term reservoir of carbon and nutrients, which can be gradually released through microbial activity.

Carbon transformation

Carbon is one of the most important elements involved in organic matter transformation. Carbon transformation is central to the entire decomposition process because carbon forms the structural framework of most organic residues and serves as an energy source for soil microorganisms. During respiration, part of this carbon is released as CO_2 . Under oxygen limited conditions, microbial processes can instead contribute to methane (CH_4) production. Another fraction of carbon becomes incorporated into microbial biomass or transformed into more persistent soil organic matter. Thus, carbon continuously moves between plant residues $\rightarrow$ microbial biomass $\rightarrow$ CO_2 $\rightarrow$ stable soil organic matter, linking decomposition with both soil fertility and the global carbon cycle. Thus, microbial succession during decomposition does not merely reduce organic residues, it progressively changes the chemical nature of organic matter and controls the timing and availability of nutrients for plants.

Factors affecting microbial succession

Microbial activity does not occur at the same rate under all conditions. Microbial succession and decomposition are influenced by a combination of environmental conditions, substrate characteristics and microbial interactions. The major factors include temperature, moisture, oxygen availability, pH, carbon-to-nitrogen (C:N) ratio, particle size and lignin content. These factors determine which microbial groups are able to grow, compete and dominate at different stages of organic matter decomposition. Temperature strongly affects microbial growth and enzyme activity. Microbial activity generally increases with temperature up to an optimum range, beyond which excessive heat can inhibit microbial growth or denature enzymes. Moisture is essential for microbial metabolism and the movement of soluble nutrients. Very dry conditions restrict microbial activity, whereas excessive moisture can fill soil pores with water and reduce oxygen availability, thereby favouring anaerobic microorganisms. Oxygen availability influences both the rate and pathway of decomposition. Well aerated conditions generally favour aerobic bacteria and fungi, whereas oxygen limited conditions favour

anaerobic microorganisms and result in different decomposition products. Soil pH also influences microbial succession because different microbial groups have different pH preferences. Changes in pH during decomposition can therefore alter the composition and activity of the microbial community.

The C:N ratio of organic residues is another important determinant of decomposition. Materials with a relatively low C:N ratio generally decompose more rapidly because microorganisms have sufficient nitrogen for growth. In contrast, residues with a high C:N ratio may decompose slowly and can promote temporary nitrogen immobilization. Particle size also affects decomposition; smaller particles provide a greater surface area for microbial colonisation and enzyme action, generally accelerating decomposition. The chemical composition and quality of the substrate further influence microbial succession. Easily degradable compounds such as sugars, amino acids and starch are utilised rapidly by early colonisers, whereas compounds such as cellulose, hemicellulose and lignin require more specialised microorganisms and enzymes. Lignin rich residues are particularly resistant to decomposition and therefore favour microbial groups capable of degrading complex plant polymers, especially certain fungi. In addition to these factors, nutrient availability, substrate composition, microbial competition and interactions among microorganisms can influence succession. As decomposition progresses, the availability of substrates changes, creating new conditions that favour different microbial groups. Thus, microbial succession is the result of a continuous interaction between the changing organic substrate and the surrounding soil environment.

Methods to study microbial succession

A combination of traditional culture based methods and modern molecular techniques is used to study microbial succession during organic matter decomposition. Traditional methods include plate counting and microscopy. In the plate count method, microorganisms are grown on suitable nutrient media and the resulting colonies are counted to estimate the abundance of culturable microorganisms. However, this method does not represent the entire microbial community because many soil microorganisms cannot be readily cultured under laboratory conditions. Microscopy provides direct visual information about microbial cells, including their size, shape and abundance, and can also help observe changes in microbial populations during decomposition. Modern molecular approaches provide a more comprehensive picture of microbial communities. DNA sequencing and metagenomics allow researchers to characterise microorganisms directly from environmental samples without the need for cultivation. These methods help to identify the microbial community by analyzing their genetic material. Quantitative PCR (qPCR) is used to quantify specific microorganisms or functional genes and thereby estimate their abundance at different stages of decomposition. 16S rRNA sequencing is mainly used to identify bacteria, while ITS (Internal Transcribed Spacer Sequencing) is used to identify fungi.

Phospholipid Fatty Acid (PLFA) analysis uses characteristic membrane lipids to estimate microbial biomass and distinguish broad microbial groups such as bacteria and fungi. Enzyme assays measure the activities of extracellular and intracellular enzymes involved in the breakdown of organic compounds and nutrient cycling, providing an indication of microbial functional activity. Metatranscriptomics analyses RNA transcripts to determine which genes are actively expressed at a particular stage, thereby providing insights into the functions being performed by the microbial community. Together, these approaches provide complementary information on the composition, diversity, abundance, biomass and functional activity of microorganisms. By comparing these parameters at different stages of decomposition, researchers can determine which microbial groups appear, increase, decline or become dominant, and how their activities contribute to the transformation of organic matter and release of nutrients.

Summary

Microbial succession is a dynamic process that drives the decomposition and transformation of soil organic matter. Different groups of microorganisms dominate successive stages, from the breakdown of simple sugars to the degradation of resistant compounds and the formation of stable humus. Through enzymatic activity, mineralisation, immobilisation and carbon transformation, these microorganisms regulate nutrient availability and contribute to long-term carbon storage. Their activities are influenced by environmental conditions and the chemical composition of organic residues. Understanding and managing this invisible microbial workforce can improve soil health, enhance nutrient cycling and support sustainable agricultural productivity.

References

1. Akanbi, E.O., Ugwuanyi, I.R., Musah, M., Quadri, O., Abu, I. and Emeribe, C. E. (2026). Influence of litter quality on microbial activity and soil nutrient cycling: A review. *Asian J. Res. Agric. For.* 12(2): 248-267.
2. Buresova, A., Kopecky, J., Hrdinkova, V., Kamenik, Z., Omelka, M. and Sagova-Mareckova, M. (2019). Succession of microbial decomposers is determined by litter type, but site conditions drive decomposition rates. *Appl. Environ. Microbiol.* 85: 24.
3. Hassan-Dalléac, S., Guiga, W. and Suau-Pernet, A. (2026). Soil microbes are the tiny bioengineers running Earth's underground factory. *Commun. Earth Environ.* 7: 403.
4. Khan, M.T., Suproniene, S., Žvirauskiene, R., and Aleinikoviene, J. (2025). Climate, Soil and Microbes: Interactions Shaping Organic Matter Decomposition in Croplands. *Agronomy* 15(8):1928.
5. Patil, R.J., Kaygude, G.A., Atole, S.K. and Muley, P.A. (2019). Role of microbes in organic matter decomposition and sustains the soil health. *Int. J. Innov. Res. Sci. Eng. Technol.* 8(7): 7756-7762.
6. Singh, R., Rani, A., Kumar, P., Shukla, G. and Kumar, A. (2016). The decomposer microorganisms in the environment and their succession of substrates. *Int. J. Eng. Sci. Res. Technol.* 1166-1171.
7. Sokol, N.W., Slessarev, E., Marschmann, G.L., Nicolas, A., Blazewicz, S.J., Brodie, E.L., Firestone, M.K., Foley, M.M., Hestrin, R., Hungate, B.A., Koch, B.J., Stone B.W., Sullivan, M.W., Zablocki, O. and Pett-Ridge, J. (2022). Life and death in the soil microbiome: how ecological processes influence biogeochemistry. *Nat. Rev.* 20: 415-429.
8. Yaseen, U., Nurbaity, A. and Fitriatin, B.N. (2026). Ecological significance of soil microbes in sustaining soil health through nutrient cycling, carbon sequestration, and structural integrity. *Discov. Soil* 3: 65.
9. Zhan, C. (2024). Microbial decomposition and soil health: mechanisms and ecological implications. *Mol. Soil Biol.* 15(2):59-70.