

Weed Management and Its Role in Reducing Crop Diseases and Insect Vectors Management

*P. G. Gameti¹ and Dr. A. R. Ninama²

¹Ph.D. Scholar (Plant Pathology), Department of Plant Pathology, B. A. College of Agriculture, Anand Agricultural University, Anand-388110, Gujarat, India

²Assistant Research scientist, Regional Research Station, Sardarkrushinagar Dantiwada Agricultural University, Bhachau, Gujarat-370140, India

*Corresponding Author's email: gametiprincy28@gmail.com

Weeds are not merely competitors for light, water, nutrients and space; some also act as alternative hosts or reservoirs for plant pathogens and insect vectors. Weeds growing within fields, on bunds, irrigation channels and field margins may allow pathogens or vectors to persist when crops are absent, potentially contributing to disease reappearance in the next crop (Wisler and Norris, 2005; Kumar *et al.*, 2021; Byron *et al.*, 2019). Their epidemiological importance is species-specific: a weed is relevant only when it supports the pathogen, vector or both under local conditions. Therefore, weed management can complement disease and insect-vector management, but indiscriminate vegetation removal is not always justified because some non-crop plants also support beneficial insects.




Figure 1. Potential pathways through which weeds can contribute to crop disease epidemics and insect-vector pressure

Weeds as Reservoirs of Plant Pathogens

Weeds can harbour viruses, bacteria, fungi and other pathogens, sometimes without conspicuous symptoms. Infected weeds may maintain inoculum between cropping periods and become sources of infection when susceptible crops are planted nearby. Weeds and volunteer

crop plants can also undermine crop-free periods intended to interrupt pathogen survival, particularly where the pathogen has a broad host range (Wisler and Norris, 2005; Byron *et al.*, 2019). In India, reviews have emphasized the indirect importance of weeds as alternate or alternative hosts for crop pests and pathogens, in addition to their direct effects on yield (Kumar *et al.*, 2021; Rao *et al.*, 2018). However, pathogen detection in a weed does not by itself prove that the weed is an epidemiologically important reservoir. Confirmation may require evidence that the pathogen remains viable and can be transmitted from the weed to the crop.

Weeds and Insect Vectors

Weeds may provide food, shelter, overwintering sites or breeding habitat for insect pests and vectors, including whiteflies, aphids, leafhoppers and thrips. When these insects acquire pathogens from infected weeds or move between weeds and crops, they can contribute to the spread of vector-transmitted diseases (Eigenbrode *et al.*, 2018; Kumar *et al.*, 2021). This is especially important for many plant viruses, for which disease risk depends on the interaction among virus, vector, host plant and landscape. Recent work on whitefly-transmitted viruses in cucurbits illustrates why field margins and non-crop vegetation should be considered during surveillance rather than focusing only on symptomatic crop plants (Weed Hosts study, 2025). Nevertheless, weed presence does not always increase vector pressure: some plants can support natural enemies or divert pests away from crops. Management decisions should therefore be based on confirmed host relationships and field observations.

Table 1. Weed-related ecological roles and their implications for crop health.

Weed-related role	How it may influence crop health	Management implication
Pathogen reservoir	Maintains some pathogens between crop seasons	Identify and target confirmed reservoir weeds
Vector host or refuge	Provides food, shelter or breeding sites for some vectors	Monitor weeds and crop margins for vectors
Volunteer crop host	Bridges pathogen or pest survival between successive crops	Remove volunteers where they are epidemiologically important
Microclimate modifier	Can alter humidity, airflow and canopy conditions	Manage dense weeds without damaging crop or soil
Beneficial-insect habitat	Some non-crop plants support predators and parasitoids	Avoid indiscriminate clearing; use selective management

How Weed Management Can Reduce Disease Risk

Effective weed management can reduce disease risk by removing confirmed pathogen reservoirs, limiting alternative hosts for vectors and reducing volunteer crops that bridge successive seasons. Priority should be given to weeds known to host the relevant pathogen or vector, particularly within the crop, along irrigation channels and at field borders. Early-season scouting and periodic monitoring of field margins help identify host weeds before vectors disperse into the crop (Byron *et al.*, 2019; Kumar *et al.*, 2021). Crop rotation, clean planting material, sanitation of tools and removal of infected volunteer plants can strengthen this approach. Weed control alone, however, rarely provides complete protection because vectors may immigrate from distant fields, pathogens may be seed-borne or soil-borne, and some vectors transmit viruses rapidly before insecticides act.

Integrated Weed and Vector Management

An integrated strategy combines cultural, mechanical, biological and chemical approaches according to crop, weed species and the target disease. Practices include competitive crop stands, mulching where suitable, crop rotation, timely hand or mechanical removal, and herbicides registered for the crop and used according to the label. Vector monitoring and management should be coordinated with weed control, because sudden destruction of heavily

infested weeds can sometimes displace mobile insects into the crop. Selective interventions and conservation of predators and parasitoids are important where beneficial organisms contribute to biological control (Eigenbrode *et al.*, 2018; Rao *et al.*, 2018). Field scouting should record weed identity, vector abundance, crop symptoms and disease incidence. Where a suspected weed reservoir is important, laboratory diagnosis and, when feasible, transmission tests can distinguish a true reservoir from a plant on which pathogen detection has little epidemiological significance.




Figure 2. A practical framework for integrated weed, disease and insect-vector management

Table 2. Integrated weed-management practices and their disease/vector-management implications.

Management practice	Potential contribution	Caution
Field and margin scouting	Finds host weeds and vector hotspots early	Identification should be accurate and repeated
Targeted weed removal	Reduces confirmed pathogen/vector reservoirs	Some mobile vectors may move into crops after removal
Crop rotation and volunteer control	Breaks host continuity for selected pests and pathogens	Check host range; broad-host-range pathogens may persist
Mechanical/cultural control	Suppresses weeds without relying solely on herbicides	Timing and crop stage affect crop safety
Label-compliant herbicides	Can provide timely control in suitable crops	Avoid off-label use, drift and herbicide-resistance selection
Vector and natural-enemy monitoring	Guides targeted action and protects biological control	Avoid unnecessary broad-spectrum insecticide use

Conclusion

Weed management is an important supporting component of integrated crop disease and insect-vector management because selected weeds can maintain pathogens, sustain vectors or connect cropping seasons. Its success depends on accurate identification of relevant weed–pathogen–vector relationships and well-timed, targeted interventions. The objective is not to eliminate all non-crop vegetation indiscriminately, but to manage confirmed reservoir weeds while protecting beneficial organisms and reducing unnecessary herbicide use. Combining weed

surveillance with crop sanitation, resistant cultivars, vector monitoring and locally validated integrated pest management offers a more durable approach than weed control alone.

References

1. Bailey, K.L., Lazarovits, G., 2003. Suppressing soil-borne diseases with residue management and organic amendments. *Soil and Tillage Research* 72(2), 169–180.
2. Blanco-Canqui, H., Ruis, S.J., 2018. No-tillage and soil physical environment. *Geoderma* 326, 164–200.
3. Bockus, W.W., Shroyer, J.P., 1998. The impact of reduced tillage on soilborne plant pathogens. *Annual Review of Phytopathology* 36, 485–500.
4. Haddaway, N.R., Hedlund, K., Jackson, L.E., Kätterer, T., Lugato, E., Thomsen, I.K., Jørgensen, H.B., Söderström, B., 2017. How does tillage intensity affect soil organic carbon? A systematic review. *Environmental Evidence* 6, 30.
5. Hobbs, P.R., Sayre, K., Gupta, R., 2008. The role of conservation agriculture in sustainable agriculture. *Philosophical Transactions of the Royal Society B: Biological Sciences* 363(1491), 543–555.
6. Jat, R.A., Sahrawat, K.L., Kassam, A. (Eds.), 2020. *Conservation Agriculture: Global Prospects and Challenges*. CABI, Wallingford.
7. Kassam, A., Friedrich, T., Derpsch, R., 2019. Global spread of conservation agriculture. *International Journal of Environmental Studies* 76(1), 29–5.
8. Pittelkow, C.M., Liang, X., Linquist, B.A., van Groenigen, K.J., Lee, J., Lundy, M.E., van Gestel, N., Six, J., Venterea, R.T., van Kessel, C., 2015. Productivity limits and potentials of the principles of conservation agriculture. *Nature* 517, 365–368.
9. Powlson, D.S., Stirling, C.M., Jat, M.L., Gerard, B.G., Palm, C.A., Sanchez, P.A., Cassman, K.G., 2014. Limited potential of no-till agriculture for climate change mitigation. *Nature Climate Change* 4, 678–683.
10. Sturz, A.V., Carter, M.R., Johnston, H.W., 1997. A review of plant disease, pathogen interactions and microbial antagonism under conservation tillage in temperate humid agriculture. *Soil and Tillage Research* 41(3–4), 169–189.