



Dry Land Integrated Farming System

*K. Malarvizhi and Keerthi D. E.

School of Agricultural Sciences, Dhanalakshmi Srinivasan University (DSU),
Samayapuram, Tiruchirappalli, Tamil Nadu, India

*Corresponding Author's email: malarvizhi3307@gmail.com

Nearly 100 million hectares of agricultural land in India depend on rainfall, making rainfed agriculture a vital contributor to the country's economy. The limited success of conventional cropping systems in improving the productivity of small farms led to the emergence of farming systems research. This approach considers the farm as an integrated system comprising crops, livestock, soil, labor, inputs, and environmental factors, where farm families strive to maximize outputs within the constraints of available resources, capabilities, and socio-cultural conditions. The adoption of integrated farming systems can enhance productivity, generate employment opportunities, increase farm income, and improve nutritional security for both people and livestock. In such systems, different components are interlinked, with the by-products or waste from one component serving as valuable inputs, food, or energy sources for others. Significant research has been undertaken to develop location-specific integrated farming systems, and a review of the available literature on integrated farming systems in dryland regions is presented below.

Characteristics of dry farming

The characteristics of dry farming in India are largely influenced by climatic conditions, particularly weather and rainfall patterns. Dryland regions account for nearly 62 percent of the country's total cultivated area of 142 million hectares and contribute about 42 percent of overall food production. These regions possess a fragile natural resource base and rely heavily on prevailing weather conditions, with rainfall being the most critical factor affecting agricultural productivity. Irregular and unevenly distributed rainfall, combined with high evaporation rates in arid climates, frequently results in water shortages that adversely affect crop stability and yields. The length of the crop-growing period in dryland areas varies considerably, ranging from 60 to 300 days, depending on factors such as total rainfall, its distribution, potential evapotranspiration, and soil moisture storage capacity. The dominant soil types in these regions are vertisols, alfisols, and their associated soil orders. These soils generally contain low levels of organic matter and are deficient in essential nutrients such as nitrogen, phosphorus, sulphur, and calcium. Moisture stress and nutrient deficiencies are the primary factors limiting crop productivity, while poor soil fertility and low water-holding capacity further contribute to reduced yields.

Dryland farming systems are often characterized by low and uncertain productivity due to inadequate utilization of rainfall and soil resources, limited application of fertilizers, insufficient adoption of high-yielding crop varieties, and poor implementation of soil conservation practices. Enhancing productivity in these regions requires efficient resource management, improved agricultural technologies, and the adoption of alternative land-use systems. Major crops cultivated under dryland conditions include sorghum, cotton, soybean, groundnut, sunflower, and various pulse crops. Crop yields in these areas are significantly affected by abiotic stresses, particularly drought, which accounts for a substantial portion of the reduction in potential productivity.

Low and highly variable rainfall remains one of the major causes of unstable production and poor yields in rainfed agriculture. The continued reliance on traditional farming methods with minimal fertilizer use has also contributed to low productivity levels. Furthermore, many rainfed ecosystems are characterized by marginal soils with poor fertility, posing additional challenges to sustainable crop production. Tamil Nadu falls within a bimodal rainfall zone, receiving an average annual rainfall of about 700 mm distributed across two rainy seasons. Water scarcity and drought are common features of the state's dryland regions. The predominant soil type is red loamy soil, accompanied by areas of sandy alluvium and red sandy soils. Red loam soils containing gravel are considered particularly suitable for livestock rearing and are relatively rich in calcium and phosphorus. Important crops grown in the dryland areas of Tamil Nadu include cotton, finger millet, and sorghum, while millet and sorghum are commonly cultivated during the monsoon season.

Integrated farming system in dry land

The success of an integrated farming system (IFS) in dryland areas depends on selecting enterprises that minimize competition for resources while maximizing complementarities among different components. Such integration enables efficient utilization of available resources, enhances productivity, and improves farm income sustainability. Studies conducted in different regions of India have demonstrated the benefits of integrating crop production with livestock enterprises. In Uttar Pradesh, farming systems combining crops, dairy, and goat rearing were found to have the highest potential, followed by crop and goat-based systems. Similarly, the integration of goat and sheep rearing with crop cultivation was recommended for the dryland regions of Punjab due to its economic viability and efficient resource utilization.

Research in Tamil Nadu identified dairy, biogas production, and silviculture as a suitable combination for garden land conditions, while crop cultivation integrated with goat rearing proved most effective under rainfed situations in the western zone of the state. In the rainfed uplands of Eastern India, farming systems that combined cereal and grain legume crops with small livestock and poultry enterprises were found to be highly beneficial. Researchers have emphasized that the successful integration of livestock within crop-based systems requires the development of appropriate, farm-oriented technologies that are socially acceptable and economically feasible. In the western zone of Tamil Nadu, dryland farming systems commonly include milch cows, goats, and buffaloes alongside crop cultivation. For a one-hectare dryland farm, an integrated model consisting of sorghum and cowpea grown for both grain and fodder, along with *Cenchrus glaucus* cultivated under *Emblica officinalis* plantations and supplemented with Tellicherry goats, was found to be more productive and profitable than sole sorghum cultivation.

Further studies recommended the integration of goats, rabbits, and pigeons with crop components such as pearl millet and soybean, maize and cowpea, and fodder grasses like *Cenchrus ciliaris* and *Stylosanthes scabra*. This diversified farming model was identified as one of the most suitable systems for one-hectare dryland farms in the western region of Tamil Nadu. An integrated farming model that included crops, goats, pigeons, buffaloes, agroforestry, and a farm pond was also evaluated in dryland conditions. In this system, Tellicherry goats were reared for meat production, buffaloes for milk, and the farm pond was used to harvest runoff water for irrigation during moisture stress periods while also serving as a source of nutrient-rich silt. The study concluded that the combination of crop cultivation enriched with composted buffalo manure, along with goats, pigeons, buffaloes, agroforestry components, and a farm pond, represented a highly suitable and sustainable farming system for the dryland regions of western Tamil Nadu. In the rainfed areas of Chhattisgarh, a farming system comprising two bullocks, one cow, one buffalo, ten goats, poultry, and ducks, integrated with crop production, was found to be particularly beneficial for marginal farmers. Similarly, in the Deogarh district of Odisha, a sustainable integrated farming system involving crops such as rice, off-season tomato, and cauliflower, combined with poultry, paddy straw mushroom cultivation, and vermicomposting, generated higher net returns and created additional employment opportunities under rainfed and risk-prone conditions.

Overall, integrated farming systems in dryland areas have proven effective in enhancing resource-use efficiency, increasing farm productivity, generating employment, improving income, and ensuring greater sustainability under challenging rainfed conditions.

Components of integrated farming system in dry land

Arable farming alone often provides only limited profits and low economic returns to farmers, making it necessary to adopt integrated farming systems that combine multiple complementary enterprises. In dryland agriculture, the integration of various farm components helps improve resource utilization, enhance productivity, diversify income sources, and reduce risks associated with climatic uncertainties. An integrated farming system may include a combination of crop production, livestock rearing, poultry, goat and sheep farming, agroforestry, horticulture, fisheries, vermicomposting, biogas production, and water-harvesting structures, depending on local resources and environmental conditions. These components interact with one another in a mutually supportive manner, where the by-products or wastes generated by one enterprise serve as valuable inputs for another. Such integration not only increases overall farm profitability but also contributes to employment generation, nutritional security, environmental sustainability, and the efficient recycling of farm resources in dryland regions.

Economic Contribution of Integrated Farming Systems in Dryland Areas

Integrated farming systems have been widely recognized for their ability to improve farm income, employment opportunities, and overall economic sustainability in dryland agriculture. Several studies have demonstrated that the inclusion of livestock enterprises, particularly buffaloes, goats, and sheep, significantly enhances the profitability of farming systems. Research has shown that buffalo rearing offers substantial economic benefits in both irrigated and rainfed regions. The integration of dairy buffaloes with existing cropping systems has been recommended as an effective strategy for improving farm income and ensuring sustainable employment generation. Studies conducted in Haryana revealed that farming systems combining crop cultivation with buffalo rearing generated higher net returns compared to systems incorporating cows or cropping alone. Similarly, the inclusion of buffaloes on small farms in Punjab was found to increase both household income and labor utilization. Small ruminants such as goats and sheep play a crucial role in the livelihoods of farmers in semi-arid regions. They represent a major source of farm income, with goat sales contributing a significant share of total household earnings. The integration of livestock with crop production also supports productivity by providing manure, utilizing crop residues as feed, and creating additional revenue streams. For example, buffaloes maintained under systems integrating sorghum and cowpea fodder crops were able to sustain satisfactory milk production levels.

Studies conducted in Tamil Nadu demonstrated the economic advantages of integrating crops, fodder grasses, fodder trees, and goat rearing. A farming system consisting of sorghum grain and fodder crops, fodder grasses, multipurpose trees, and Tellicherry goats generated substantially higher income than crop cultivation alone. Goat rearing accounted for the majority of the total income from the system, while the integrated approach also created considerable additional employment opportunities throughout the year. The importance of goats and sheep in mixed farming systems has been highlighted across Asia, where they occupy a valuable economic and ecological niche. In the dryland regions of southern Tamil Nadu, integrating crop production with rainfed fruit trees and goat rearing significantly increased the income of small and marginal farmers. Likewise, the introduction of sheep into rubber plantations was reported to enhance farm profitability considerably by providing an additional source of revenue without adversely affecting plantation productivity. Economic evaluations of various enterprise combinations have further confirmed the profitability of livestock integration. Farming systems combining crops such as wheat and sugarcane with buffalo rearing generated high net returns, while integrated models involving sugarcane cultivation, dairying, and sheep rearing produced even greater profits. These findings indicate that diversified farming enterprises are generally more profitable than reliance on crop production alone.

Silvipasture-based farming systems have also demonstrated strong economic potential in dryland environments. Among livestock enterprises, goat rearing generated the highest income, followed by dairy cattle. Similarly, integrated crop–tree–goat systems in the western region of Tamil Nadu produced substantially higher net returns compared to sole cropping systems under rainfed conditions.

In the rainfed black soil regions of southern Tamil Nadu, the integration of tree legumes such as *Leucaena leucocephala*, *Acacia senegal*, and *Prosopis cineraria* with perennial fodder grasses and goat rearing provided farmers with a significant additional annual income from relatively small landholdings. These results emphasize that integrated farming systems not only improve profitability but also enhance livelihood security and economic resilience in dryland agriculture.

Overall, the economic benefits of integrated farming systems arise from diversification, efficient resource utilization, reduced production risks, increased employment generation, and the creation of multiple income sources, making them a sustainable and profitable approach for dryland farming regions.

Employment Generation through Integrated Farming Systems in Dryland Areas

Integrated farming systems (IFS) play a significant role in generating employment opportunities in dryland regions, where agricultural activities are often seasonal and farm families experience unemployment for a large part of the year. Since crop cultivation in rainfed areas is generally confined to the monsoon season, integrating additional enterprises such as livestock, poultry, agroforestry, biogas production, and mushroom cultivation helps ensure year-round employment and income generation. Studies have shown that integrated farming systems create substantially more employment than crop production alone. For instance, sole crop cultivation generated about 400 man-days of employment, whereas an integrated farming system that included six buffaloes produced as many as 904 man-days annually. Similarly, goat rearing has been recognized as an effective means of employment generation, particularly in dryland areas where alternative livelihood opportunities are limited.

The integration of sheep and goat enterprises within farming systems has been found to be especially beneficial because these animals can utilize natural grazing resources while requiring relatively low investment. Research has indicated that large-scale goat rearing generates the highest labour employment among livestock enterprises, particularly under grazing-based management systems. In addition to contributing to farm income, sheep and goat husbandry provides regular and continuous employment to farm households throughout the year. Studies conducted in Uttar Pradesh revealed that crop farming, buffalo rearing, goat rearing, and sheep farming contributed significantly to household labour utilization, with crop and buffalo enterprises accounting for the largest share of employment. Likewise, in the arid regions of Rajasthan, the adoption of silvipastoral and hortipastoral systems integrated with sheep and goat rearing created substantial additional employment opportunities for rural households. Research in Tamil Nadu demonstrated that integrated farming systems involving crops and goats generated an additional 314 man-days of employment per hectare annually under rainfed conditions. Furthermore, systems integrating crop production with dairy farming, biogas units, and mushroom cultivation provided even greater employment, generating up to 875 man-days per year. These diversified enterprises not only increased labour requirements but also ensured a steady distribution of work throughout the year.

Dryland farming models incorporating sorghum, cowpea, fodder grasses, fodder trees, and goat rearing have also been shown to generate significant additional employment compared to conventional cropping systems. Similarly, integrated systems combining crops, dairy farming, biogas production, silviculture, and spawn production created several hundred additional man-days of employment annually over cropping alone in irrigated garden lands. Goat farming, in particular, has proven to be a major source of rural employment. It has been reported that maintaining a herd of approximately 200 goats can provide full-time employment

for at least two individuals throughout the year. Under dryland conditions, both sheep and goat rearing have consistently demonstrated their ability to generate employment while remaining economically profitable, especially when animals are managed through grazing-based systems. Overall, integrated farming systems contribute significantly to employment generation by diversifying farm enterprises, reducing seasonal unemployment, improving labour utilization, and creating sustainable livelihood opportunities for rural households in dryland regions. Such systems are particularly valuable for small and marginal farmers, as they provide continuous work and enhance economic stability throughout the year.

Resource Recycling in Integrated Farming Systems

Resource recycling is a fundamental feature of integrated farming systems, particularly in smallholder agriculture, where crop and livestock enterprises interact closely and share available natural resources. The efficient recycling of farm residues and wastes helps improve soil fertility, reduce production costs, and enhance the sustainability of farming systems. Studies have shown that integrated crop–livestock systems minimize nutrient losses by ensuring that nutrients removed from the field through fodder harvest are returned in the form of animal manure. Research conducted in Tamil Nadu demonstrated that the application of goat manure to crop fields improved both soil fertility and crop productivity. Such recycling practices contribute to maintaining soil health while reducing dependence on chemical fertilizers. The integration of crops and livestock facilitates the effective utilization of agricultural wastes and by-products. Livestock consume crop residues and fodder produced on the farm, while the manure generated by animals is recycled back to the soil as an organic nutrient source. This continuous nutrient cycling enriches soil organic matter, improves soil structure, and enhances nutrient availability for crops. Studies in dryland and rainfed regions have emphasized that the recycling of livestock wastes significantly contributes to nutrient conservation and organic enrichment of soils.

Animal manure serves as an important resource for maintaining soil fertility. It has been reported that a pair of draught animals can produce substantial quantities of manure annually, which can be applied to crop fields to sustain soil productivity. Integrated farming systems therefore reduce farmers' reliance on externally purchased inputs by utilizing resources already available within the farm. Researchers have highlighted that integrated farming systems offer excellent opportunities for recycling crop residues, fodder, animal wastes, and other farm by-products. Crop residues are used as livestock feed, while livestock wastes are converted into valuable organic manure for crop cultivation. This approach promotes efficient resource utilization and contributes to environmental sustainability. Studies conducted in the western region of Tamil Nadu found that the combined application of chemical fertilizers and goat manure improved soil fertility and enhanced nutrient recycling efficiency. The integration of dairy and goat enterprises with crop production further strengthened resource utilization, as crop residues served as feed for animals, and the nutrients removed through harvest were returned to the soil through livestock manure.

More diversified integrated farming systems involving crops, pigeons, goats, buffaloes, agroforestry components, and farm ponds have been shown to generate larger quantities of organic residues and improve waste recycling efficiency. Such systems produce significant amounts of organic biomass that can be converted into nutrient-rich manure, thereby improving nutrient availability and soil productivity. Research has also indicated that integrated farming systems generate higher quantities of residues with greater nutrient value compared to sole cropping systems. The incorporation of livestock and agroforestry components increases the availability of organic materials that can be recycled within the farm. Composting and vermicomposting of crop residues and animal manure further enhance nutrient recovery and provide valuable organic fertilizers for field crops, fodder crops, and vegetables. The integration of vegetable cultivation, food crops, goat rearing, and vermicomposting has been found to improve overall farm productivity under irrigated upland conditions. Recycling vermicompost and goat manure not only reduces the cost of cultivation by lowering fertilizer requirements but also improves soil health, enhances microbial activity, and supports

sustainable crop production. Overall, resource recycling in integrated farming systems promotes efficient utilization of farm resources, conserves nutrients, improves soil fertility, reduces dependence on external inputs, and enhances the long-term sustainability and profitability of both dryland and irrigated agricultural systems.

Conclusion

A review of the existing farming systems revealed that the various farm enterprises often function independently, with inadequate linkages among system components. The introduction of appropriate technological interventions and the integration of complementary enterprises have been shown to improve the overall performance of farming systems and enhance the economic well-being of farm families. Integrated farming systems have proven to be more profitable, resilient, and sustainable than conventional monocropping systems. The evaluation of different enterprise combinations through physical indicators provides valuable insights into system performance and resource-use efficiency. Indicators such as employment generation, nutritional contribution, energy efficiency, resource utilization, and residue recycling help in identifying and optimizing suitable combinations of farm enterprises.

The integration of crops, livestock, agroforestry, and other complementary components promotes efficient utilization of available bioresources, enhances nutrient recycling, and reduces wastage within the farming system. In addition, these systems contribute significantly to employment generation and provide diverse sources of income, thereby improving livelihood security for farming households. Furthermore, integrated farming systems play an important role in ensuring nutritional security by supplying a variety of food products for both humans and livestock. Their ability to maximize resource recycling, improve productivity, strengthen economic returns, and provide sustainable employment opportunities makes them a highly suitable approach for the dryland farming regions. Therefore, the adoption of integrated farming systems can serve as an effective strategy for achieving sustainable agricultural development and improving the socio-economic conditions of farmers in dryland areas.

References

1. Acharya, S.S., K.A. Varghese and Diwakar Singh. (1987). Economic aspects of dairy enterprise in an irrigated arid region of Rajasthan. *Indian J. Dairy Sci.*, 40(3): 331–338.
2. Aggarwal, R.K. and P. Kumar. (1993). Sustainable productivity through organic manure in arid land ecosystem. In: *Proceedings of International Symposium on Environmental Degradation*, CAZRI, Jodhpur. p. 21.
3. Alam, M.R., R.I. Sarker, M.D. Hossain and M.S. Islam. (2000). Contribution of livestock to small farms in Bangladesh. *Asian Australasian J. Anim. Sci.*, 13(suppl.): 339–342.
4. Arvind, R. and K.K. Jain. (1992). Dairying potential in Jalandhar district of Punjab. *Indian Dairyman*, 44(12): 578–583.
5. Barik, K.C., D. Mohanty, S.K. Nath, S.K. Sahoo and Laba Soren. (2010). Assessment of integrated farming system for rainfed agro ecosystem in Deogarh district of Orissa. In: *Extended summaries of XIX National Symposium on Resource Management Approaches towards Livelihood Security*, Bengaluru, Karnataka, Dec. 2–4. p. 172.
6. Basavaraj, H. and N.R. Gangadharappa. (1999). Adoption level and profitability of different combination of farm enterprises in existing farming systems. *Mysore J. Agric. Sci.*, 33(2): 261–266.
7. Das, S.K. (2001). Prospect and potentiality for goat farming in north eastern region of India – a review. *Agric. Rev.*, 22(3/4): 228–233.
8. Deoghare, P.R. and N.K. Bhattacharya. (1993). Economic analysis of goat rearing in the Mathura district of Uttar Pradesh. *Indian J. Anim. Sci.*, 64(12): 1368–1372.
9. Deoghare, P.R. (1997). Sustainability of on-farm income and employment through livestock production in Mathura district. *Indian J. Anim. Sci.*, 67(16): 916–919.
10. Devendra, C. (1998). Improvement of small ruminant production systems in rainfed agro-ecological zones of Asia. *Ann. Arid Zone*, 37(3): 215–232.

11. Devendra, C. (2000). Animal production and rainfed agriculture in Asia: Potential opportunities for productivity enhancement. *Outlook on Agriculture*, 29: 161–175.
12. Esther Shekinah, D., C. Jayanthi and N. Sankaran. (2005). Physical indicators of sustainability – A farming systems approach for the small farmer in the rainfed vertisols of the Western zone of Tamil Nadu. *Journal of Sustainable Agriculture*, 25(3): 43–65.
13. Gajja, B.L., T.K. Bhati, L.N. Harsh and M.S. Khan. (1999). Comparative economics of silvipasture, hortipasture and annual crops on marginal agricultural lands of arid zone of Rajasthan. *Ann. Arid Zone*, 38(2): 173–180.
14. Gupta, P.C., K. Singh and M.A. Akbar. (1994). Developing forage based rations for lactating buffaloes. In: *Proceedings of IV World Buffalo Congress*, Sao Paulo, Brazil, June 27–30, Vol. II. pp. 275–277.
15. Jayanthi, C., M. Balusamy, C. Chinnusamy and S. Mythili. (2003). Integrated nutrient supply system of linked components in lowland integrated farming system. *Indian J. Agron.*, 48: 241–246.
16. Jayanthi, C., C. Vennila, K. Nalini and G. Vivek. (2007). Physical indicators – on farm farming system for irrigated uplands of Western zone of Tamil Nadu. *Journal of Farming Systems Research and Development*, 13(1): 17–25.
17. Kadian, V.S., K.P. Singh, S.N. Singh and H. Kumar. (1992). Productivity and economics of intensive crop rotations in arable and mixed farming systems. *Haryana J. Agron.*, 8(2): 116–122.
18. Kale, A. (1999). Goat farming – a remunerative enterprise. *Intensive Agriculture*, 37(3&4): 24–28.
19. Nehra, O.P., N. Singh, S.N. Singh, V. Kumar, K.P. Singh and V.S. Kadian. (1992). Comparative studies on economics of various farming systems under semi-arid tracts of Southern Haryana. *Indian J. Anim. Prod. Mgmt.*, 8: 121–124.
20. Pandey, R.N. (1988). Contribution of buffaloes to generation of income and employment for the rural households, especially for the small farmers. *Indian J. Anim. Prod. Mgmt.*, 4(3/4): 124–132.
21. Pandey, R.N. and T.S. Bhogal. (1980). Prospects of increasing income and employment on mixed farms. *Indian J. Agric. Econ.*, 35(4): 144–151.
22. Pathak, P. and K.B. Laryea. (1995). Soil and water conservation in the Indian SAT; principles and improved practices. In: R.P. Singh (Ed.), *Sustainable Development of Dryland Agriculture in India*. Scientific Publishers, Jodhpur. pp. 83–92.
23. Phengvichith, V. (1998). The integration of livestock in shifting cultivation systems in the upland of the Lao PDR. *FFTC Extension Bulletin*, 464: 1–10.
24. Policy Note. (2010). Agriculture Department. Government of Tamil Nadu.
25. Radhamani, S. (2001). Sustainable integrated farming system for dryland Vertisol areas of Western Zone of Tamil Nadu. Ph.D. Thesis, Tamil Nadu Agricultural University, Coimbatore.
26. Rai, K.N. and S.N. Singh. (1982). Income and employment effects of livestock and new dryland farm technology in Haryana. *Indian J. Dairy Sci.*, 35: 203–208.
27. Ramasamy, C., S. Natarajan, C. Jayanthi and D. Suresh Kumar. (2007). Intensive integrated farming system to boost income of farmers. In: *Proceedings of 32nd IAUA Vice Chancellors Annual Convention on Diversification in Indian Agriculture*, Birsa Agricultural University, Dec. 20–21. pp. 28–47.
28. Ramrao, W.Y., S.P. Tiwari and P. Singh. (2006). Crop-livestock integrated farming system for marginal farmers in rainfed regions of Chhattisgarh in Central India. *Livestock Research for Rural Development*, 18(7): 23–30.
29. Rangasamy, A. (1995). Integrated farming systems research in Tamil Nadu. Country paper on development of sustainable farming systems. National Productivity Organization, Sri Lanka, June 5–10.

30. Rangasamy, A., V.S. Shanmugasundaram, S. Sankaran and M. Subbarayalu. (1990). Integrated farming system management: A viable approach. Research Bulletin, Tamil Nadu Agricultural University, Coimbatore.
31. Reddy, S.R. (2000). *Principles of Agronomy*. Kalyani Publishers, New Delhi. pp. 149–179.
32. Sajise, P.E. (1998). Ecological concerns in crop-livestock integration in sloping land. *FFTC Extension Bulletin*, 461.
33. San Nu Nu and B.J. Deaton. (1999). Feasibility of integrating sheep and crops with small holder rubber productions in Indonesia. *Journal of Agribusiness*, 17(2): 105–122.
34. Senthilvel, T., K.R. Latha and N. Gopalasamy. (1998). Farming system approach for sustainable yield and income under rainfed vertisols. *Madras Agric. J.*, 55(1): 65–67.
35. Sheshshayee, M.S., H. Bindumadhava, A.G. Shanker, T.G. Prasad and M. Udayakumar. (2003). Breeding strategies to explain water use efficiency for crop improvement. *Journal of Plant Biology*, 30(2): 253–268.