



Green Barriers, Healthy Crops: An Eco-Friendly Way to Manage Insect-Vector-Viral Diseases

*Swati Saha

ICAR–Indian Agricultural Research Institute, Regional Centre, Pune, India

Corresponding Author's email: swatisaha1980@gmail.com

Vegetable cultivation plays a vital role in Indian agriculture by providing essential nutrients, generating employment and improving farmers' incomes. However, viral diseases remain a major constraint to vegetable production, often causing substantial losses in yield, quality and marketability. Diseases such as tomato leaf curl, chilli leaf curl and papaya ringspot can seriously affect crop growth and productivity. Many plant viruses are transmitted by insects such as aphids, whiteflies and thrips. Although insecticides are commonly used to control these vectors, they are not always effective in preventing virus transmission. In particular, viruses transmitted in a non-persistent manner can be acquired and transmitted by aphids within a short period during probing. The virus may therefore spread before an insecticide can kill the vector. Repeated pesticide applications also increase production costs, encourage insecticide resistance, harm beneficial insects and contribute to environmental pollution and pesticide residues in produce. These challenges highlight the need for preventive, economical and environmentally friendly disease management strategies. Border cropping offers one such practical approach by creating a protective green barrier around the main crop.

What Is Border Cropping?

Border cropping involves growing selected plants along the boundaries of a field, or in strategically arranged rows within a field, to reduce pest movement and protect the main crop. These plants are often called barrier crops or protector crops because they can interfere with the movement and feeding behaviour of insect vectors. The technique is a form of habitat manipulation that makes the cropping environment less favourable for pests while supporting a more balanced agricultural ecosystem. Suitable barrier crops include maize, sorghum and pigeon pea (arhar). Depending on the crop, local conditions and the insect vector involved, these plants can act as physical barriers, influence insect landing and movement, and reduce the number of vectors reaching the main crop. The effectiveness of a barrier crop depends on its height, density, arrangement, suitability for the local farming system and relationship with the target insect and virus. Ideally, it should not serve as a reservoir host for the virus or its vector.

How Does the Green Barrier Work?

Border cropping can protect crops through several complementary mechanisms.

1. Intercepting incoming insect vectors: A sufficiently tall and dense border may interfere with the movement of flying insects into the main crop. It can reduce direct entry and alter the flight paths of some vectors, although the degree of protection varies with insect behaviour and field conditions.

2. Disrupting virus transmission: Aphids transmitting non-persistent viruses may briefly probe a plant and acquire or transmit virus particles during feeding. When these aphids land on a non-host barrier crop, repeated probing may cause them to lose their ability to transmit certain viruses. If they subsequently move to the main crop, the likelihood of successful virus transmission may be reduced. This effect depends on the particular virus–vector relationship.

Therefore, barrier crops must be selected carefully rather than assuming that every border plant will prevent virus spread.

3. Influencing insect landing and feeding: The physical characteristics of barrier crops, including leaf texture and plant structure, can influence insect landing and feeding behaviour. Some insects may be less inclined to settle and feed on certain non-host plants, while others may land on them temporarily. This can help disrupt their movement towards the susceptible crop.

4. Supporting integrated pest management: When combined with regular field monitoring, removal of infected plants where appropriate, weed-host management and selective insecticides or biopesticides, border cropping can contribute to an integrated approach to pest and virus management. It is important to remember that barrier crops do not provide complete protection against all insect vectors or viral diseases. Their main role is to reduce risk as part of a broader preventive strategy.

Establishing Border Crops: A Practical Guide for Farmers

Correct timing and arrangement are essential for obtaining the best results from border cropping.

Step 1: Select a suitable barrier crop. Choose crops such as maize, sorghum or pigeon pea according to the main vegetable crop, the target insect vector, local growing conditions and available space. The selected barrier should preferably be a non-host of the target virus and vector.

Step 2: Sow the barrier in advance. Border crops may be sown approximately 30–45 days before planting the main crop, depending on the species and local growing conditions. This allows the barrier to establish sufficient height and density before the vegetable crop becomes vulnerable to incoming vectors.

Step 3: Maintain a dense and continuous border. Sow the barrier rows closely enough to create an effective boundary, while following suitable spacing and agronomic recommendations for the selected crop. Gaps in the border may allow insects to enter the main field more easily.

Step 4: Consider single or double borders. A single border of maize, sorghum or pigeon pea may be suitable under some conditions. Where additional protection is needed, a combination of two barrier crops may be considered, provided the arrangement is agronomically suitable and does not create alternative pest or virus reservoirs.

Step 5: Monitor the main crop regularly. Inspect plants for symptoms of viral infection and monitor the presence of insect vectors. Remove infected plants where recommended and practical, particularly during the early stages of crop growth, when virus infection can cause severe yield losses.

Step 6: Use pesticides judiciously. If vector populations or other pests require intervention, use locally recommended, selective insecticides or biopesticides as part of integrated pest management. Border cropping should complement, rather than automatically replace, other necessary control measures.

Benefits of Border Cropping

1. Reduced dependence on chemical pesticides: By helping to limit the movement of certain insect vectors, border crops may reduce the need for repeated insecticide applications. This can lower input costs and help slow the development of pesticide resistance when combined with responsible pesticide use.

2. Protection of beneficial insects: Excessive pesticide use can harm natural enemies such as ladybird beetles, lacewings and parasitoids. Reducing unnecessary sprays helps conserve these beneficial organisms and supports natural pest regulation.

3. Healthier crops and better productivity: Reducing early virus infection can help preserve plant growth, yield and produce quality. The benefits will depend on the disease, vector pressure, crop susceptibility and effectiveness of the barrier arrangement.

4. Environmental protection and safer produce: Fewer unnecessary pesticide applications can reduce environmental contamination and the risk of pesticide residues on harvested vegetables. Farmers must still observe all applicable label directions and pre-harvest intervals whenever pesticides are used.

5. Potential economic benefits: Barrier crops may reduce expenditure on pesticides and, in some systems, provide an additional harvest or other farm benefits. However, their establishment requires land, seed, labour and management. Farmers should consider these costs and the possible competition for water, nutrients and sunlight when planning the border.

6. Greater sustainability in vegetable production: Border cropping is a practical cultural component of integrated pest management. It encourages preventive crop protection, reduces exclusive dependence on chemical control and can contribute to more resilient farming systems.

Where Can This Technique Be Used?

Border cropping may be useful in vegetable production systems affected by insect-vectored viral diseases, including tomato, chilli, cucurbits and other susceptible crops. However, the choice of barrier crop must be matched to the target virus and vector. For example, an approach intended to reduce aphid-mediated transmission of a non-persistent virus may not provide the same level of protection against a virus transmitted by whiteflies. Similarly, a barrier that works under one set of climatic and cropping conditions may perform differently elsewhere. Therefore, farmers should follow locally validated recommendations from agricultural universities, ICAR institutes and extension specialists. Demonstrations and field trials can help identify suitable barrier crops, planting dates and row arrangements for individual farming situations.

Conclusion: Let Nature Help Protect the Crop

Border cropping demonstrates how a simple change in field design can contribute to more sustainable crop protection. By planting suitable barrier crops around vegetable fields, farmers can interfere with the movement of certain insect vectors and potentially reduce the spread of viral diseases. Although this method cannot eliminate virus infection or replace every other control measure, it offers an environmentally friendly addition to integrated pest management. When combined with healthy planting material, timely crop establishment, regular monitoring, removal of infected plants where appropriate and judicious pesticide use, border cropping can help reduce avoidable crop losses and dependence on chemical inputs.

A green barrier is more than a row of plants—it is a step towards healthier crops, safer food and sustainable farming.



Field layout with double border crop (Arhar and Maize)



Okra and Chilli crop planted with maize as border crop for management of Okra Yellow Vein mosaic virus and Leaf curl virus